

# In vitro effects of oxytocin, acepromazine, detomidine, xylazine, butorphanol, terbutaline, isoproterenol, and dantrolene on smooth and skeletal muscles of the equine esophagus

Anne A. Wooldridge, DVM, MS; Susan C. Eades, DVM, PhD; Giselle L. Hosgood, BVSc, MS; Rustin M. Moore, DVM, PhD

**Objective**—To characterize the in vitro effects of oxytocin, acepromazine, xylazine, butorphanol, detomidine, dantrolene, isoproterenol, and terbutaline on skeletal and smooth muscle from the equine esophagus.

**Animals**—14 adult horses without digestive tract disease.

**Procedure**—Circular and longitudinal strips from the skeletal and smooth muscle of the esophagus were suspended in tissue baths, connected to force-displacement transducers interfaced with a physiograph, and electrical field stimulation was applied. Cumulative concentration-response curves were generated for oxytocin, acepromazine, xylazine, detomidine, butorphanol, isoproterenol, terbutaline, and dantrolene. Mean maximum twitch amplitude for 3 contractions/min was recorded and compared with predrug-vehicle values for the skeletal muscle segments, and area under the curve (AUC) for 3 contractions/min was compared with predrug-vehicle values for the smooth muscle segments.

**Results**—No drugs caused a significant change in skeletal muscle response. In smooth muscle, isoproterenol, terbutaline, and oxytocin significantly reduced AUC in a concentration-dependent manner. Maximum reduction in AUC was 69% at  $10^{-4}$ M for isoproterenol, 63% at  $10^{-5}$ M for terbutaline, and 64% at  $10^{-4}$ M for oxytocin.

**Conclusions and Clinical Relevance**—Isoproterenol, terbutaline, and oxytocin cause relaxation of the smooth muscle portion of the esophagus. The clinical relaxant effects on the proximal portion of the esophagus reported of drugs such as oxytocin, detomidine, and acepromazine may be the result of centrally mediated mechanisms. (*Am J Vet Res* 2002;63:1732–1737)

Esophageal obstruction is the most commonly diagnosed esophageal disorder of horses.<sup>1</sup> Xylazine, detomidine, acepromazine, butorphanol, and oxytocin are used in the treatment of this condition to decrease pain and anxiety, lower the horse's head, and potentially relax the spasm around the obstruction.<sup>1,2</sup> The effects of these drugs on equine esophageal muscle have not been critically elucidated in vitro, and there are few evaluations in vivo.<sup>3-5</sup>

The equine esophagus is composed of skeletal muscle in the proximal two-thirds and smooth muscle in the distal one-third.<sup>1,6</sup> Most obstructions occur in the skeletal muscle portion at the post-pharyngeal area or at the area of the thoracic inlet.<sup>1</sup> Denac et al<sup>2</sup> evaluated the effect of various neurotransmitters on smooth muscle preparations of equine esophagus in vitro. Norepinephrine caused concentration-dependent relaxation of longitudinal esophageal smooth muscle from the esophageal body via its agonist effects on  $\beta$  receptors.<sup>2</sup> Acetylcholine and norepinephrine (via  $\alpha$  receptors) caused concentration-dependent contraction of the circular smooth muscle of the esophageal body and the lower esophageal sphincter.<sup>2</sup> Electrical field stimulation (EFS) was also applied to the smooth muscle preparations, and the effects were characterized.<sup>2</sup> The skeletal muscle portion of the equine esophagus has not been tested in vitro.

We hypothesized that xylazine, detomidine, oxytocin, butorphanol, acepromazine, and dantrolene would cause relaxation of skeletal esophageal muscle, and isoproterenol and terbutaline would have no effect on skeletal muscle. We also hypothesized that isoproterenol, terbutaline, xylazine, detomidine, acepromazine, and butorphanol would cause relaxation of esophageal smooth muscle, dantrolene would have no effect, and oxytocin would have either no effect or cause contraction. The purpose of the study reported here was to characterize the effects of these drugs on isolated preparations of skeletal and smooth muscle from the equine esophagus.

## Materials and Methods

**Tissue harvest and preparation**—This study was approved by the Louisiana State University Institutional Animal Care and Use Committee. Muscle from the esophageal body was harvested from 14 adult horses that were euthanatized for reasons unrelated to digestive tract disease. Tissue was collected immediately after euthanasia with an overdose of pentobarbital sodium<sup>a</sup> (100 mg/kg, IV). Skeletal esophageal tissue was collected from the cer-

Received Jan 9, 2002.

Accepted Jun 4, 2002.

From the Equine Health Studies Program, Department of Veterinary Clinical Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, LA 70803. Dr. Wooldridge's present address is the Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC 27710.

This report represents a portion of the thesis submitted by the first author to the graduate school, Louisiana State University, in partial fulfillment of the requirements for the MS degree.

Presented at the American College of Veterinary Internal Medicine Forum, Chicago, June 1999, and the 7th International Colic Research Symposium, Manchester, England, July 2002.

The authors thank Catherine Koch and Michael Broussare for technical assistance.

Address correspondence to Dr. Wooldridge.

vical region, excluding the upper esophageal sphincter, and smooth muscle was collected from the distal thoracic region, excluding the lower esophageal sphincter. The tissue was immediately placed into warmed, oxygenated Krebs solution and transported to the laboratory. The Krebs solution composition was NaCl (118mM), KCl (4.7mM), NaHCO<sub>3</sub> (84.01 g/L), MgSO<sub>4</sub> (1.2mM), KH<sub>2</sub>PO<sub>4</sub> (1.1mM), CaCl<sub>2</sub> × 2 H<sub>2</sub>O (2.5mM), and 3.5 g/L of dextrose. The esophagus was opened longitudinally, and the mucosa and submucosa were gently removed using sharp dissection, leaving the muscular layers. The muscle was oriented so that 2 circular and then 2 longitudinal muscle strips, 1.5 × 0.5 cm, were collected from each of the skeletal muscle and smooth muscle portions of the esophagus. The 8 preparations were suspended in individual tissue baths containing 14 mL of Krebs solution, maintained at 37°C, and continuously aerated with 95% O<sub>2</sub> and 5% CO<sub>2</sub>. One end of each strip was fixed with silk suture to a force-displacement transducer<sup>b</sup> interfaced with a polygraph<sup>c</sup>, and the other end was fixed to the base of the tissue bath.

**Electrical field stimulation**—Electrical field stimulation was accomplished by suspending each tissue preparation between 2 platinum wire electrodes mounted on the tissue bath. A square wave electrical impulse was generated by a stimulator<sup>d</sup> and applied to all 8 tissues. Repetitive stimuli called trains (60 V, 5 Hz, 0.5 ms for skeletal muscle, and 90 V, 10 Hz, 1 ms for smooth muscle) were applied at a rate of 0.025 trains/s. Pilot studies in our laboratory determined that these parameters resulted in submaximal stimulation for these skeletal and smooth muscle tissues as reported for equine esophageal smooth muscle and rabbit urethral skeletal muscle.<sup>2,7</sup>

**Experimental design**—Tissues from 3 horses were evaluated without EFS, and tissues from 11 horses were evaluated with EFS. The experiments on unstimulated tissues were performed to evaluate spontaneous activity. Electrical field stimulation was used for more consistent contractions and because a relaxation response could be easily identified and accurately measured.

For evaluation of tissue responses without EFS, smooth and skeletal muscle tissues were allowed to equilibrate in the baths for 1 hour, and then resting tension was set to 1 to 2 g. Tissues were washed with fresh Krebs solution at 15-minute intervals, and the tension was reset to 1 to 2 g at each washing until the final washing before drug administration. The tissues were first observed for spontaneous activity for 15 minutes. A pilot study was performed on the smooth and skeletal muscle to determine a drug-induction of maximum contraction amplitude. The drugs tested in the pilot study were histamine, neostigmine, norepinephrine, oxytocin, and carbachol. Oxytocin was included with the other drugs because it was expected to cause contraction of smooth muscle, and we wanted to determine its effect on unstimulated esophageal skeletal muscle, given the recent interest in the drug. Cumulative concentrations of histamine<sup>e</sup> (10<sup>-9</sup>, 10<sup>-8</sup>, 10<sup>-7</sup>, 10<sup>-6</sup>, 10<sup>-5</sup>, 10<sup>-4</sup>M), KCl<sup>f</sup> (10<sup>-4</sup>M), neostigmine<sup>g</sup> (10<sup>-9</sup>, 10<sup>-8</sup>, 10<sup>-7</sup>, 10<sup>-6</sup>, 10<sup>-5</sup>M), BaCl<sub>2</sub><sup>h</sup> (10<sup>-3</sup>M), norepinephrine<sup>i</sup> (10<sup>-11</sup>, 10<sup>-10</sup>, 10<sup>-9</sup>, 10<sup>-8</sup>, 10<sup>-7</sup>, 10<sup>-6</sup>, 10<sup>-5</sup>M), oxytocin<sup>j</sup> (10<sup>-7</sup>M), and carbachol<sup>k</sup> (10<sup>9</sup>, 10<sup>-8</sup>, 10<sup>-7</sup>, 10<sup>-6</sup>, 10<sup>-5</sup>M) were individually added to the tissue baths to reach the desired concentration in the bath. Tissues were washed with fresh Krebs solution between drugs. Carbachol was the only drug to have any effect, and it caused a concentration-dependent increase in tension in both smooth and skeletal muscle, with maximum contraction amplitude first occurring at 10<sup>-6</sup>M. After determining a drug-induction of maximum contraction amplitude in the pilot study, an experiment was performed to evaluate a relaxant effect of oxytocin after precontraction with carbachol

(10<sup>-6</sup>M). After equilibration and establishment of resting baseline of the smooth and skeletal muscle tissues, tension was increased by placing carbachol (10<sup>-6</sup>M) into the bath. When the response plateaued, oxytocin (10<sup>-12</sup>, 10<sup>-11</sup>, 10<sup>-10</sup>, 10<sup>-9</sup>, 10<sup>-8</sup>, 10<sup>-7</sup>M) or vehicle (water) was added to the tissues to determine a cumulative concentration-response relationship. Parameters measured included baseline, contraction amplitude, and frequency before carbachol administration, after carbachol administration, and for each concentration of oxytocin or vehicle.

Smooth and skeletal muscle tissues from 11 horses were used for EFS. After preparation, the tissues were allowed to equilibrate for 1 hour, and then resting tension was set to 1 to 2 g. Tissues were washed with fresh Krebs solution at 15-minute intervals throughout the experiment. Electrical field stimulation was applied for 15 minutes, and the tissues were washed and rested for another 15 minutes. The drugs tested on the stimulated tissues included oxytocin (10<sup>-12</sup> to 10<sup>-4</sup>M), acepromazine<sup>l</sup> (10<sup>-9</sup> to 10<sup>-4</sup>M), isoproterenol<sup>m</sup> (10<sup>-9</sup> to 10<sup>-4</sup>M), terbutaline<sup>n</sup> (10<sup>-9</sup> to 10<sup>-3</sup>M), xylazine<sup>o</sup> (10<sup>-9</sup> to 10<sup>-4</sup>M), butorphanol<sup>p</sup> (10<sup>-9</sup> to 10<sup>-4</sup>M), detomidine<sup>q</sup> (10<sup>-9</sup> to 10<sup>-4</sup>M), dantrolene<sup>r</sup> (10<sup>-9</sup> to 10<sup>-3</sup>M), and vehicle control (water). For oxytocin, only lower concentrations (10<sup>-12</sup> to 10<sup>-7</sup>M) were used for the first 2 horses, and more concentrations (10<sup>-6</sup>, 10<sup>-5</sup>, and 10<sup>-4</sup>M) were added for the next 9 horses. This resulted in analysis of more tissues for the lower concentrations of oxytocin in the final analysis. Drugs or vehicle were administered at 1-minute intervals in increasing concentrations; 10 μL of each drug dilution or 10 μL of vehicle (water) was added to the baths to achieve the final drug concentration in the bath. Electrical field stimulated-contractions were recorded for 1 minute before drug administration. Contractions were measured for 1 minute after each concentration of drug or vehicle was added to the baths. Acepromazine was always administered last because of the need for acid-washing the tissue baths after use of this drug. The other drugs or vehicle was administered in random order. Baths were rinsed extensively with Krebs solution 3 times after the highest concentration of each drug was administered, and the tissues were rested for 15 minutes between drugs. Tetrodotoxin<sup>s</sup> (3 × 10<sup>-7</sup>M), which blocks sodium channels so that a nerve-stimulated action potential cannot be generated, was administered after the final washing to determine that responses to EFS were nerve-mediated.

For skeletal muscle, stimulation caused 3 sets of twitches/min. Maximum twitch amplitudes (grams) of the 3 sets were measured with calipers, and the mean was calculated for 1 minute before drug or vehicle administration and for each drug or vehicle concentration. Percentage change from mean maximum twitch amplitude before drug administration was calculated for each drug concentration or vehicle.

For smooth muscle, stimulation caused 3 sustained contractions (rather than sets of twitches) per minute. **Area under the curve**<sup>t</sup> (AUC) was the measurement performed on contractions to account for duration as well as maximum contraction amplitude. This was measured and averaged for 3 contractions for 1 minute before drug or vehicle administration and for each drug concentration or vehicle control. Percentage change from mean AUC before drug or vehicle administration was calculated for each drug concentration or vehicle.

For stimulated smooth and skeletal muscle, tissues with mean maximum contraction or twitch amplitudes before drug or vehicle administration of < 0.5 g were excluded from the experiment because they were considered to be nonviable tissues.

**Statistical analyses**—The data were found to follow a normal distribution after arcsine transformation by use of

the Shapiro-Wilk statistic, with failure to reject the null hypothesis of normality at  $P < 0.05$ . The transformed data were analyzed by use of a mixed effect general linear model procedure, PROC MIXED,<sup>8,9</sup> which accounted for the random variance of horse and repeated measurements on each horse. There was no difference in response of longitudinal versus circular layers, so the data were pooled and reanalyzed with the general linear model procedure. When there were significant effects of drugs at  $P < 0.05$ , predetermined comparisons from mean maximum twitch amplitude or AUC before drug or vehicle administration and from vehicle controls were made by use of the differences of least squares means. For all comparisons, a value of  $P < 0.05$  was considered significant. A significant percentage of change attributable to a drug was considered to occur if there was a significant difference from percentage change before drug or vehicle administration and from the vehicle percentage change occurring at the same time interval as that concentration of drug. Both of these comparisons were made to determine a difference from values before drug or vehicle administration and to control for tissue fatigue over time. All data are reported as mean  $\pm$  SEM.

## Results

**Unstimulated tissues**—Analysis of unstimulated tissues was performed on 4 tissues for oxytocin for smooth muscle and 4 for vehicle control (4 were nonviable), and on 5 tissues for oxytocin for skeletal muscle and 4 for vehicle control (3 were nonviable). The tissues tended to relax after the resting tension was set, and the baseline would return to 0 g of tension. Smooth muscle had spontaneous activity, whereas skeletal muscle did not. Carbachol ( $10^{-6}$ M) raised the baseline tension, increased frequency of contractions, and increased amplitude of contractions in smooth and skeletal muscle. Baseline tension in skeletal muscle increased from resting baseline of 0.0 g to stimulated baseline of  $0.8 \pm 0.2$  g after administration of carbachol. Carbachol increased the maximum contraction amplitude from 0 to  $4.8 \pm 2.0$  g in skeletal muscle. The frequency of spontaneous contractions in skeletal muscle increased from 0.0 to  $4.5 \pm 0.4$ /min after carbachol administration. In smooth muscle, baseline tension increased from  $0.25 \pm 0.07$  g (resting baseline) to  $0.95 \pm 0.23$  g (stimulated baseline) after carbachol administration. Carbachol increased the maximum contraction amplitude from  $0.55 \pm 0.17$  to  $4.2 \pm 0.78$  g in smooth muscle. The frequency of contraction in the smooth muscle increased from  $1.75 \pm 0.53$  to  $3 \pm 0.35$ /min after carbachol administration.

There was no difference between tissues precontracted with carbachol and treated with oxytocin versus those treated with a vehicle in either smooth or skeletal muscles.

**Stimulated skeletal muscle strips**—Thirty tissues were analyzed for each drug or vehicle for skeletal muscle (except for the lower concentrations of oxytocin, which used 20 tissues). Fourteen tissues were not included in analysis because they were nonviable. There was no significant response to any of the tested drugs and no evidence of tissue fatigue over time (Fig 1). Tetrodotoxin decreased the

response to stimulation, confirming nerve-mediated responses to EFS.

**Stimulated smooth muscle strips**—Twenty-three smooth muscle strips were analyzed for each drug or vehicle for the stimulated muscle experiment (except for the higher concentrations of oxytocin, which used 16). Twenty-one tissues were nonviable. Tetrodotoxin decreased the response to stimulation, confirming a nerve-mediated response to EFS. The AUC for the vehicle-treated tissues at each interval corresponding to increasing concentrations of the drug were all significantly lower than the AUC before vehicle administration, indicating tissue fatigue across time. Isoproterenol, terbutaline, and oxytocin caused a significant decrease in AUC when compared with AUC before drug administration and with the vehicle percentage change corresponding to that concentration of drug, indicating a significant effect of drug. Isoproterenol caused a significant decrease in AUC at concentrations of  $10^{-6}$ M,  $10^{-3}$ M, and  $10^{-4}$ M, and terbutaline caused a decrease in AUC at  $10^{-5}$ M (Fig 1 and 2). The maximum decrease in AUC for isoproterenol was 69% at  $10^{-4}$ M, and for terbutaline was 63% at  $10^{-5}$ M. Oxytocin caused a decrease in AUC at  $10^{-8}$ ,  $10^{-5}$ , and  $10^{-4}$ M, and the maximum decrease was 64% at  $10^{-4}$ M (Fig 3). Xylazine, acepromazine, butorphanol, detomidine, and dantrolene did not cause a significant decrease in AUC at any drug concentration.




Figure 1—A—Response (maximum twitch amplitude [g]) of electrical field stimulated (EFS) skeletal esophageal muscle to increasing concentrations of oxytocin. Notice no change in amplitude in response to oxytocin (Oxy) and no tissue fatigue over time. B—Response (area under the curve [AUC]) of EFS esophageal smooth muscle to increasing concentrations of isoproterenol (Iso). Notice a decrease in AUC in response to isoproterenol.




Figure 2—Mean percentage change in AUC in EFS esophageal smooth muscle preparations ( $n = 23/\text{group}$ ) treated with isoproterenol, terbutaline, or vehicle, in response to increasing drug or vehicle concentrations. \*Significant ( $P < 0.05$ ) difference from AUC before drug administration and from AUC of tissues exposed to vehicle alone.




Figure 3—Mean percentage change in AUC in EFS esophageal smooth muscle preparations ( $n = 23/\text{group}$  unless otherwise indicated) treated with oxytocin or vehicle, in response to increasing drug or vehicle concentrations. \*Significant ( $P < 0.05$ ) difference from AUC before drug administration and from AUC of tissues exposed to vehicle alone.

## Discussion

No drugs significantly affected esophageal skeletal muscle in our model, but several drugs caused a decrease in AUC in the smooth muscle. We evaluated drug-induced muscle contraction and EFS to find the best method to study consistent contraction and subsequent drug-induced relaxation. Carbachol stimulates G-protein coupled muscarinic receptors, resulting in muscle contraction. It is not metabolized by acetylcholinesterase as quickly as is acetylcholine. Carbachol did induce increases in baseline and maximum contraction amplitude, and it also increased spontaneous contractions by changing the transmembrane potential. The EFS-tissues had more consistent contractions and were considered to be a better model for studying potential relaxant effects of the tested drugs.

We hypothesized that oxytocin, detomidine, xylazine, butorphanol, acepromazine, and dantrolene would cause a decrease in maximum twitch amplitude

in skeletal esophageal muscle. We expected oxytocin, a peptide hypothalamic hormone, to cause a decrease in maximum twitch amplitude in the skeletal muscle, because it was reported in 2 *in vivo* studies that oxytocin caused a pressure decrease and clinical resolution of luminal obstructions in the proximal esophagus.<sup>3,9</sup> Because oxytocin did not cause a decrease in maximum amplitude in our model *in vitro*, the effect observed *in vivo* could be due to a centrally mediated mechanism rather than a local mechanism. The smooth muscle relaxation caused by oxytocin could potentially cause the skeletal muscle to relieve pressure as well as to open the passageway for food boluses to pass. Oxytocin has been implicated as a neurotransmitter or neuromodulating agent and affects nociception in the spinal cord.<sup>10</sup> Esophageal skeletal muscle activity is controlled by nerve-mediated stimulation from the brain, causing stimulation of nicotinic receptors via the vagus nerve.<sup>11</sup> Oxytocin could be acting as a neuromodulator of the central nerve-stimulated activity of the proximal portion of the esophagus *in vivo*.

Dantrolene was expected to cause a decrease in maximum twitch amplitude in the skeletal muscle of the esophagus *in vitro*, because it is a direct-acting skeletal muscle relaxant that interferes with excitation-contraction coupling at the sarcoplasmic reticulum and at the T-tubule system by changing calcium release and reuptake.<sup>12</sup> The effects of dantrolene on the equine esophagus have not been studied, but have been studied *in vitro* on sections of skeletal muscle from the external urethral sphincter in rats and humans.<sup>7</sup> Dantrolene did cause a decrease in nerve-mediated (tetrodotoxin sensitive) twitch amplitude *in vitro* in the external urethral sphincter and had no effect on the smooth muscle of the bladder in rats.<sup>7</sup> Dantrolene had no effect on the skeletal muscle in our study, possibly because the effect on skeletal esophageal muscle is different from that on the external urethral sphincter, or because conditions (eg, pH, calcium concentration) in the bath may not have been optimal for an effect. Because skeletal muscle in the external urethral sphincter is interlaced with smooth muscle and the innervation to that area is complex, responses to drugs could be quite different from responses in esophageal skeletal muscle.<sup>7</sup>

Detomidine and xylazine,  $\alpha$ -2 receptor agonists, had no effect on skeletal muscle *in vitro*. Detomidine has profound inhibitory effects on motility of the proximal portion of the esophagus in horses when studied with contrast radiography *in vivo*.<sup>5,13</sup> Neither of these drugs has been evaluated on skeletal muscle of the esophagus *in vitro*, but a nonspecific  $\alpha$ -receptor blocking agent, phentolamine, has been evaluated on the external urethral sphincter *in vitro* in rats and had no effect, suggesting that local  $\alpha$ -receptors are not important in the stimulation or relaxation of that skeletal muscle *in vitro*.<sup>7</sup> The effects of detomidine observed with contrast radiography *in vivo* may be through a central, rather than a locally, mediated mechanism, which explains the lack of response in the skeletal muscle seen *in vitro*. Neither butorphanol, an opiate agonist-antagonist, nor acepromazine, a phenothiazine tranquilizer, has been reported to cause much effect on

esophageal motility in vivo, but are reported to be clinically useful.<sup>1,13,14</sup> Neither drug caused any changes in the skeletal muscle in vitro, so the effects observed clinically may be from pain and anxiety relief rather than from direct relaxation of the esophageal muscle.

The smooth muscle of the equine esophagus is arranged in circular and longitudinal layers that contain an intrinsic nerve plexus, which plays a role in regulation of motility.<sup>2,6</sup> Smooth muscle contraction activity is controlled by nerve-mediated stimulation from the plexus of nerves in the esophageal wall.<sup>11,15</sup> Our in vitro study concentrated on drugs that affect the muscular response, although certain intact organs in a live animal could have different responses to these drugs because the submucosa, which also contains neuronal cell bodies that transmit to the myenteric plexus, would be intact. The myenteric plexus was also disturbed when circular muscle sections were taken; nevertheless, there was no significant difference between circular and longitudinal muscle responses, so the data were pooled.

Acetylcholine is the major stimulating neurotransmitter in smooth muscle of the equine esophagus.<sup>2</sup> Norepinephrine caused contraction of the lower esophageal sphincter that was fully blocked by prazosin, an  $\alpha$ -1 receptor blocking agent.<sup>2</sup> Norepinephrine caused relaxation of the longitudinal muscle of the esophageal body that was fully blocked by propranolol, a nonspecific  $\beta$ -receptor blocking agent.<sup>2</sup> This would indicate that  $\beta$ -receptors are important for relaxation of the smooth muscle of the esophageal body. As would be expected, isoproterenol (a nonspecific  $\beta$ -receptor agonist) and terbutaline (a  $\beta$ -2 receptor agonist) caused a significant decrease in AUC in the smooth muscle in this model.

Oxytocin induces profound myometrial contraction near the time of parturition. It also causes contraction of the acini of the mammary gland causing milk let-down.<sup>16</sup> This effect may be due to an increase in calcium influx through receptor-operated channels and release of intracellular calcium.<sup>17</sup> Oxytocin also causes contraction of the aorta, basilar arteries, and cremaster muscle of rats; carotid arteries of dogs; and urinary bladder of rabbits.<sup>17-21</sup> However, oxytocin also causes vasodilatation in pulmonary vasculature of rats and cerebral arteries in dogs and rats.<sup>22-24</sup> This vasodilatory response is probably associated with nitric oxide release or V1-vasopressinergic receptor stimulation.<sup>22-24</sup> Oxytocin caused a decrease in tension in the smooth muscle in our study, which could have contributed to the beneficial effects of oxytocin in the treatment of proximal obstructions observed in vivo, potentially by opening the distal passageway for the obstruction to pass, which lessens the workload of the proximal portion of the esophagus.<sup>3,8</sup> A mechanism of action was not elucidated, but may be mediated by nitric oxide release or the V1-vasopressinergic receptor stimulation, as in some vessels.

No drugs tested in vitro had a significant effect on the proximal portion of the esophagus. The clinical effects of drugs such as oxytocin, detomidine, xylazine, acepromazine, and butorphanol may be attributable to centrally mediated relaxation of the esophagus. The sedatives and analgesics may also be causing relief of pain and anxiety rather than indirectly relaxing esophageal spasm.

The relaxant effect of isoproterenol, terbutaline, and oxytocin on smooth muscle could be clinically useful with distal obstructions, although most obstructions occur in the proximal portion of the equine esophagus.

<sup>a</sup>Beuthanasia-D Special, Schering-Plough Animal Health Corp, Kenilworth, NJ.

<sup>b</sup>Grass F-D transducer, Grass Instruments Co, Quincy, Mass.

<sup>c</sup>Grass model 79 physiograph, Grass Instruments Co, Quincy, Mass.

<sup>d</sup>Grass S48 stimulator, Astro-Med Inc, Grass Instruments Division, West Warwick, RI.

<sup>e</sup>Histamine, Sigma-Aldrich Inc, St Louis, Mo.

<sup>f</sup>KCl, Sigma-Aldrich Inc, St Louis, Mo.

<sup>g</sup>Neostigmine, Sigma-Aldrich Inc, St Louis, Mo.

<sup>h</sup>BaCl<sub>2</sub>, Sigma-Aldrich Inc, St Louis, Mo.

<sup>i</sup>Norepinephrine, Sigma-Aldrich Inc, St Louis, Mo.

<sup>j</sup>Oxytocin, Sigma-Aldrich Inc, St Louis, Mo.

<sup>k</sup>Carbachol, Sigma-Aldrich Inc, St Louis, Mo.

<sup>l</sup>Acepromazine maleate, Fort Dodge Laboratories, Fort Dodge, Iowa.

<sup>m</sup>Isoproterenol bitartrate salt, Sigma-Aldrich Inc, St Louis, Mo.

<sup>n</sup>Terbutaline sulfate, Novartis Pharmaceuticals Corp, East Hanover, NJ.

<sup>o</sup>Xylazine, Sigma-Aldrich Inc, St Louis, Mo.

<sup>p</sup>Detomidine hydrochloride, Pfizer Animal Health, West Chester, Pa.

<sup>q</sup>Butorphanol tartrate, Fort Dodge Laboratories, Fort Dodge, Iowa.

<sup>r</sup>Dantrolene sodium, Procter and Gamble Pharmaceuticals, Cincinnati, Ohio.

<sup>s</sup>Tetrodotoxin, Sigma-Aldrich Inc, St Louis, Mo.

<sup>t</sup>Sigma Scan Pro 4.0, Jandel Scientific Software, San Rafael, Calif.

<sup>u</sup>SAS V.8.0, SAS Institute, Cary, NC.

## References

1. Green EM, MacFadden KW. Esophageal disorders of the horse. In: Smith BP, ed. *Large animal internal medicine*. 2nd ed. St Louis: Mosby Year Book Inc, 1996;698-709.
2. Denac M, Bebie P, Scharer E. Effect of various neurotransmitters and electrical field stimulation on smooth muscle preparations from the esophagus of horses. *Zentralbl Veterinarmed A* 1993;40:501-508.
3. Hance SR, Noble J, Holcombe S, et al. Treating choke with oxytocin, in *Proceedings*. Am Assoc Equine Pract 1997;338-339.
4. Meyer GA, Helms RJ, Rashmir-Raven A, et al. Effect of oxytocin on contractility of the equine esophagus: treatment for esophageal obstruction, in *Proceedings*. Am Assoc Equine Pract 1997;336-337.
5. Watson TD, Sullivan M. Effects of detomidine on equine oesophageal function as studied by contrast radiography. *Vet Rec* 1991;129:67-69.
6. Slocombe RF, Todhunter RJ, Stick JA. Quantitative ultrastructural anatomy of esophagus in different regions in the horse: effects of alternate methods of tissue processing. *Am J Vet Res* 1982;3:1137-1142.
7. Massimo P, Manzini S. The rat external urethral sphincter: an in vitro model to evaluate the activity of drugs on the smooth and striated components of the urinary bladder outlet. *J Pharmacol Toxicol Methods* 1992;28:8590.
8. *SAS user's guide: statistics, version 8 edition*. Cary, NC: SAS Institute Inc, 1999.
9. Meyer G, Rashmir-Raven A, Helms R, et al. The effect of oxytocin on contractility of the equine oesophagus: a potential treatment for oesophageal obstruction. *Equine Vet J* 2000;32:151-155.
10. Brown DC, Perkowski S. Oxytocin content of the cerebrospinal fluid of dogs and its relationship to pain induced by spinal cord compression. *Vet Surg* 1998;27:607-611.
11. Weisbrodt N. Swallowing. In: Johnson L, ed. *Gastrointestinal physiology*. 5th ed. St Louis: Mosby Year Book Inc, 1997;23-31.
12. Morgan K, Bryant S. The mechanism of action of dantrolene sodium. *J Pharmacol Exp Ther* 1977;201:138-147.
13. King JN, Davies JV, Gerring EL. Contrast radiography of the equine oesophagus: effect of spasmolytic agents and passage of a nasogastric tube. *Equine Vet J* 1990;22:133-135.

14. Gaynor F, Hoffer RE, Nichols MF. Physiologic features of the canine esophagus: effects of tranquilization on esophageal motility. *Am J Vet Res* 1980;41:727-732.
15. Weisbrodt N. Regulation: nerves and smooth muscle. In: Johnson L, ed. *Gastrointestinal physiology*. 5th ed. St Louis: Mosby Year Book Co 1997;16-21.
16. Al-EknaH M, Homeida A. A review of some aspects of pharmacology of oxytocin in domestic animals. *Vet Res Commun* 1991;15:45-55.
17. Barrigon S, Tamargo J. The effect of oxytocin on contractile responses and <sup>45</sup>Ca movements in rat isolated aortic strips. *Br J Pharmacol* 1986;87:763-770.
18. D'Orleans-Juste P, Dion S, Mizrahi J. Effects of peptides and non-peptides on isolated arterial smooth muscles: role of endothelium. *Eur J Pharmacol* 1985;114:9-21.
19. Szabo C. Regional differences in the contractile activity of neuropeptide Y, endothelin, oxytocin, and vasopressin: comparison with non-peptidergic constrictors. An in vitro study in the basilar and mesenteric arteries of the rat. *Acta Physiol Hung* 1992;79:281-293.
20. Romine M, Anderson G. Evidence for oxytocin receptors in the urinary bladder of the rabbit. *Can J Physiol Pharmacol* 1985;63:287-291.
21. Elliott A. The effect of oxytocin on blood vessels of cremaster muscle of the rat. *Br J Pharmacol* 1975;53:149-151.
22. Kadekaro M, Liu H, Terrell M. Role of NO on vasopressin and oxytocin release and blood pressure responses during osmotic stimulation in rats. *Am J Physiol* 1997;273:1024-1030.
23. Russ R, Resta T, Walker B. Pulmonary vasodilatory response to neurohypophyseal peptides in the rat. *J Appl Physiol* 1992;73:473-478.
24. Oyama H, Suzuki Y, Satoh S, et al. Role of nitric oxide in the cerebral vasodilatory responses to vasopressin and oxytocin in dogs. *J Cereb Blood Flow Metab* 1993;13:285-290.