

8. Breig A, Turnbull I, Hassler O. Effects of mechanical stresses on the spinal cord in cervical spondylosis. A study on fresh cadaver material. *J Neurosurg* 1966;25:45–56.
9. Breig A, El-Nadi AF. Biomechanics of the cervical spinal cord. Relief of contact pressure on and overstretching of the spinal cord. *Acta Radiol Diagn (Stockh)* 1966;4:602–624.
10. Breig A. Overstretching of and circumscribed pathological tension in the spinal cord—a basic cause of symptoms in cord disorders. *J Biomech* 1970;3:7–9.
11. Breig A, Troup JDG. Biomechanical considerations in the straight-leg-raising test. Cadaveric and clinical studies of the effects of medial hip rotation. *Spine* 1979;4:242–250.
12. Yamada S, Won D, Kido D. Adult tethered cord syndrome: new classification correlated with symptomatology, imaging and pathophysiology. *Neurosurg Q* 2001;11:260–275.
13. Fletcher TF. Spinal cord and meninges. In: Evans HE, ed. *Miller's anatomy of the dog*. Philadelphia: WB Saunders Co, 1993;800–828.
14. Blumfeld H. Spinal nerve roots. In: Blumfeld H, ed. *Neuroanatomy through clinical cases*. Sunderland, Mass: Sinauer Associates Inc, 2002;303–338.
15. Sugawara O, Atsuta Y, Iwahara T, et al. The effects of mechanical compression and hypoxia on nerve root and dorsal root ganglia. An analysis of ectopic firing using an in vitro model. *Spine* 1996;21:2089–2094.
16. Kobayashi S, Yoshizawa H, Nakai S. Experimental study on the dynamics of lumbosacral nerve root circulation. *Spine* 2000;25:298–305.
17. Kobayashi S, Shizu N, Suzuki Y, et al. Changes in nerve root motion and intradiscal blood flow during an intraoperative straight-leg-raising test. *Spine* 2003;28:1427–1434.
18. Fahrni WH. Observations on straight leg-raising with special reference to nerve root adhesions. *Can J Surg* 1966;9:44–48.
19. Kjellby-Wendt G, Carlsson SG, Styf J. Results of early active rehabilitation 5–7 years after surgical treatment for lumbar disc herniation. *J Spinal Disord Tech* 2002;15:404–409.
20. Kjellby-Wendt G, Styf J, Carlsson SG. Early active rehabilitation after surgery for lumbar disc herniation: a prospective, randomized study of psychometric assessment in 50 patients. *Acta Orthop Scand* 2001;72:518–524.
21. Kjellby-Wendt G, Styf J. Early active training after lumbar discectomy. A prospective, randomized, and controlled study. *Spine* 1998;23:2345–2351.
22. Danielsen JM, Johnsen R, Kibsgaard SK, et al. Early aggressive exercise for postoperative rehabilitation after discectomy. *Spine* 2000;25:1015–1020.
23. Carragee EJ, Helms E, O'Sullivan GS. Are postoperative activity restrictions necessary after posterior lumbar discectomy? A prospective study of outcomes in 50 consecutive cases. *Spine* 1996;21:1893–1897.
24. Scrimshaw S, Maher C. Randomized controlled trial of neural mobilization after spinal surgery. *Spine* 2001;26:2647–2652.



**Correction: Neurohormonal, hemodynamic, and electrocardiographic evaluations of healthy dogs receiving long-term administration of doxorubicin**

In “Neurohormonal, hemodynamic, and electrocardiographic evaluations of healthy dogs receiving long-term administration of doxorubicin” (*AJVR*, Vol 67, pp 1319–1325), the corresponding author, Dr. Rute Chamié Alves de Souza’s address should read, “Av. Visconde de Albuquerque, 258, apto.603, 50610-090, Recife-PE, Brazil.”