

Characterization of the responses of equine digital veins and arteries to calcitonin gene-related peptide

Lisa M. Katz, DVM, PhD; Celia M. Marr, BVMS, MVM, PhD; Jonathan Elliott, VetMB, PhD

Objective—To compare responses of equine digital arteries (EDAs) and veins (EDVs) to human- α calcitonin gene-related peptide (h α CGRP), evaluate effect of the endothelium, and characterize receptors and sources of endogenous CGRP.

Sample—Palmar digital vessels (5 to 9/experiment) from healthy adult horses killed at an abattoir.

Procedures—Vessel rings were mounted under tension in organ baths containing Krebs-Henseleit solution at 30°C, with relaxation responses examined in vessels precontracted with a thromboxane-mimetic (3×10^{-8} M). Responses of endothelium-intact (+e) and -denuded (-e) EDAs and EDVs to h α CGRP (10^{-10} to 3×10^{-7} M) were compared. Following incubation with an h α CGRP receptor antagonist (h α CGRP₈₋₃₇; 1 μ M), responses of EDA(-e) and EDV(-e) to h α CGRP (10^{-7} M) were obtained. Responses of endothelium-intact and -denuded arteries and veins to h α CGRP (3×10^{-7} M) or capsaicin (10^{-5} M) were evaluated as well as responses of endothelium-intact and -denuded EDA and EDV to h α CGRP (10^{-10} to 10^{-8} M) after incubation with endothelin-1 (ET-1; 10^{-12} M).

Results—h α CGRP resulted in nonendothelium, concentration-dependent relaxation in EDAs and EDVs, with greater responses in EDAs. Treatment with h α CGRP₈₋₃₇ had minimal effect on responses to h α CGRP in either vessel type. Capsaicin induced relaxation in both vessel types. There were no differences between responses to h α CGRP for vessels pretreated with ET-1 or vehicle.

Conclusions and Clinical Relevance—Both h α CGRP and capsaicin induced digital vasodilation unaffected by a functional endothelium. This suggested that endogenous CGRP likely emanates from sensory-motor nerves and may contribute to digital vasodilation. (*Am J Vet Res* 2011;72:975-981)

Digital vasoconstriction leading to increased postcapillary resistance has been proposed to be the initiating factor that leads to clinical signs of acute laminitis in horses.^{1,2} Because vascular tone in small blood

Received May 29, 2010.

Accepted June 2, 2010.

From the University Veterinary Hospital, School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland (Katz); the Department of Veterinary Clinical Sciences, The Royal Veterinary College, University of London, North Mymms, Hatfield Hertfordshire, AL9 7TA, England (Marr); and the Department of Veterinary Basic Sciences, The Royal Veterinary College, University of London, Royal College St, London NW1 0TU, England (Elliott). Dr. Marr's present address is Rossdales Equine Hospital & Diagnostic Centre, Cotton End Rd, Exning, Newmarket, Suffolk, CB8 7NN, England.

Supported by a Clinical Research Fellowship from The Royal Veterinary College and the President's Research Award from the University College Dublin.

Presented in part at the 19th Annual Forum of the American College of Veterinary Internal Medicine, Denver, May 2001, and at the 46th British Equine Veterinary Association Congress, Edinburgh, September 2007.

The authors thank David Byrne, Erin Bennett, Turlough McNally, and Teresa Quinn for technical assistance.

Address correspondence to Dr. Katz (lisa.katz@ucd.ie).

ABBREVIATIONS

CGRP	Calcitonin gene-related peptide
CRC	Concentration response curve
DMSO	Dimethyl sulfoxide
EDA	Equine digital artery
EDA(+e)	Endothelium-intact equine digital artery
EDA(-e)	Endothelium-denuded equine digital artery
EDV	Equine digital vein
EDV(+e)	Endothelium-intact equine digital vein
EDV(-e)	Endothelium-denuded equine digital vein
ET-1	Endothelin-1
h α CGRP	Human- α calcitonin gene-related peptide
NO	Nitric oxide
pA ₂	Negative logarithm of the molar concentration of an antagonist that reduces the effect of a dose of agonist to that of half the dose
SP	Substance P

vessels is under neuronal and endothelial cell control,³ dysfunction of either may contribute to the pathogenesis of acute laminitis. It has been proposed by some researchers that peripheral neuropathy contributing to

an alteration of digital blood flow, similar to that observed in human diabetic patients⁴ or in humans with Raynaud's phenomenon,⁵ could contribute to the development of laminitis in horses. The pathogenesis of Raynaud's phenomenon is unknown, but it has been hypothesized that sensory-motor nerve dysfunction involving depletion of vasodilatory neuropeptides, such as CGRP, contributes to the development of the disease.^{3,6,7} Interestingly, investigators in another study⁸ detected abnormal morphology of peripheral nerves as well as a reduced number of unmyelinated and myelinated nerve fibers in the digital nerves of horses with laminitis, compared with results for clinically normal horses.

Calcitonin gene-related peptide is part of the calcitonin family and is involved in nociception, neurogenic inflammation, neurotransmission, and regulation of vascular tone.⁹⁻¹⁴ In rats and humans, 2 main isoforms of CGRP have been identified as α CGRP (CGRP-I) and β CGRP (CGRP-II),¹⁵ whereas more recently, the cDNA encoding for equine CGRP-I and -II has been cloned and sequenced.¹⁶ Both CGRP-I and CGRP-II are synthesized in the CNS and peripheral nervous system and have similar biological activities in most vascular beds.^{15,17} However, CGRP-I predominates in sensory pathways¹⁵ and is considered the most potent and abundant vasodilatory neuropeptide identified.^{18,19} It is commonly found in unmyelinated sensory afferent fibers and is usually colocalized with SP.²⁰⁻²² Its relaxant effects can be mediated by direct action on vascular smooth muscle receptors or indirectly through the release of endothelial-derived relaxing factors²³ (eg, NO); these relaxant effects differ between vascular beds.²⁴

It has been proven in an immunohistochemical study²⁵ that CGRP and SP often coexist in the same perivascular terminal. In equine tissues, CGRP has been identified with SP in the respiratory tract,²⁶⁻³⁰ reproductive tract,³¹ synovial joints,^{32,33} and feet.^{34,35} More specifically to equine digits, CGRP-like immunoreactivity has been detected within the hoof dermis,³² navicular suspensory apparatus, and distal interphalangeal joint.³⁵ In addition, CGRP-containing sensory-motor nerves innervating arteries and arteriovenous anastomoses have been identified within equine feet,³⁶ although their physiologic importance remains to be determined. Receptors for CGRP have been identified within the media and intima of resistance vessels in several species,³⁷ although to our knowledge, no CGRP-like sensory receptor has been identified in association with CGRP-like immunoreactivity in equine digits, which leads to the suggestion that the CGRP-containing nerve fibers are naked nerve endings.³² Only 1 published study³⁸ has been conducted to evaluate functional responses of equine tissues to CGRP. In that study,³⁸ investigators found that exogenous CGRP had no effect on tracheal smooth muscle.

Because it has been determined that SP has an endothelial-dependent relaxant effect on both EDAs and EDVs³⁹ and is often colocalized with CGRP, the examination of equine digital vascular responses to CGRP is fundamental to understanding its physiologic importance as a potential counter-regulatory vasodilatory mediator in equine digits. We hypothesized that CGRP

could be involved in vascular regulation in equine digits. Therefore, the objectives of the study reported here were to examine the relative potency and efficacy of CGRP as a vasodilator of EDVs and EDAs obtained from clinically normal horses, establish the effect of the endothelium on these responses, determine whether the observed responses were receptor-mediated events, and evaluate a possible source for endogenous CGRP in the equine digital vasculature.

Materials and Methods

Sample—Digital arteries and veins were collected from the forelimbs of the cadavers of mixed-breed healthy adult horses (males and females) killed at an abattoir. Horses were stunned by use of a captive bolt, which was followed by exsanguination; this was conducted under the supervision of a veterinarian and in accordance with European Union regulations. Forelimbs were removed within 10 minutes after stunning and exsanguination, and the limbs and feet were examined. Any limbs with gross pathological lesions were rejected from the study. For each forelimb, the digital artery was cannulated at the level of the metacarpophalangeal joint, and 180 mL of ice-cold modified Krebs-Henseleit solution (ie, Krebs solution) was infused through a 16-gauge, 52-mm catheter.^a The skin was then reflected to expose the digital artery and coronary venous plexus, which were dissected free, placed in ice-cold Krebs solution, and transported to our laboratory. On arrival at our laboratory, connective tissue was removed from the vasculature, and the arteries and veins were cut into rings of 3 to 4 mm in length. Vessel rings that were not used immediately were stored overnight in oxygenated Krebs solution at 4°C.

Experimental procedure—When necessary, the endothelium was removed by gently rubbing the intimal surface of a vessel with a wooden cocktail stick⁴⁰; this technique has been confirmed microscopically and functionally^{41,42} to be an appropriate method. Vessel rings were placed between 2 parallel wires within an organ bath containing Krebs solution maintained at 30°C and aerated with 95% O₂ and 5% CO₂. One wire was fixed in place and the other connected to an isometric force transducer,^b with the signal sent to a 2-channel pen recorder^c via a bridge amplifier.^d Four experiments were conducted, and for each experiment, a basal resting tension of 3 and 2 g was applied to arteries and veins, respectively; tissues were allowed to equilibrate under resting tension for a period of at least 1 hour. These resting tensions had previously been assessed to be optimal for mimicking *in vivo* diastolic vascular tone and allowing the greatest degree of contraction to be obtained.^{43,44} Six vessel rings could be examined at 1 time, with the vessels paired from the same horse when direct comparison was required within a given experiment. Only 1 cumulative CRC was obtained for each vessel ring. Experiments in which antagonists were tested were performed by the use of adjacent vessel rings from the same horse for the control and antagonist-treated vessels.

All vessels were constricted initially with depolarizing Krebs solution (118mM KCl) to evaluate

smooth muscle function; vessels were discarded if they constricted to < 50% of their resting tension. After peak tension was attained, vessels were washed with Krebs solution to return the muscle to the basal resting tension. Following return to basal resting tension, a thromboxane-mimetic (U44069^e [9, 11-dideoxy-9 α , 11 α -epoxymethano-prostaglandin F_{2 α} ; 3 \times 10⁻⁸M]) was added to the organ bath to provide a stable increase in vessel tension against which vasodilators could be assessed; if the vessel's resultant contractile response to the thromboxane mimetic was < 50% of its response to the depolarizing Krebs solution, then the vessel was discarded. Once the increase in tension had stabilized, a cholinergic agonist (carbachol^e) was added cumulatively (10⁻⁸ to 10⁻⁵M for experiments 1 and 2) or as a single concentration (1 μ M for experiments 3 and 4) to confirm the presence or absence of a functional endothelium. We considered a functional or intact endothelium to be present for the EDAs and EDVs if the U44069-induced tension was changed by at least 40% and 50%, respectively, in response to 1 μ M carbachol. The endothelium was considered adequately denuded if the vessels relaxed < 10% of the U44069-induced tension. The concentration of U44069 used in the study reported here had been previously determined to induce approximately 80% of the maximum increase in tension induced by U44069 in equine tissue of this size.⁴¹ The concentration of carbachol used (1 μ M) in the study reported here had also been previously found to cause maximal inhibition of the U44069-induced tension in this size and type of tissue⁴¹; further increases in the concentration of carbachol had been found to cause an increase in tension in some tissues.⁴¹ All vessels were then washed with drug-free Krebs solution to return them to basal resting tension. When applicable, antagonists or the vehicle control treatment used in the experiment was added to the organ baths at this point, and tissues were allowed to incubate for 30 minutes before addition of the vasoconstrictor.

EXPERIMENT 1—DIGITAL VASCULAR RESPONSES TO CGRP AND EFFECT OF THE ENDOTHELIUM ON THESE RESPONSES

Following return of the vessel rings to basal resting tension, U44069 (3 \times 10⁻⁸M) was added to the bathing solution to increase tension. Once a steady tension was established, CRCs to h α CGRP^e (10⁻¹⁰ to 10⁻⁶M) were obtained for paired endothelium-intact and -denuded EDA and EDV from 6 horses. The decrease in tension was expressed as a percentage of the U44069-induced tension in that same vessel ring and plotted against the log₁₀ CGRP concentration.

EXPERIMENT 2—EFFECT OF A PEPTIDE CGRP-RECEPTOR ANTAGONIST ON EDA(-E) AND EDV(-E) RESPONSES TO CGRP

Responses of EDA(-e)s (n = 5; 1/horse) or EDV(-e)s (n = 5; 1/horse) to a single concentration of h α CGRP (10⁻⁷M) were obtained in vessels precontracted by the addition of U44069 (3 \times 10⁻⁸M) before and after a 30-minute incubation with a peptide CGRP-receptor antagonist (h α CGRP₈₋₃₇^e; 1 μ M) or the same volume of 1% acetic acid (vehicle control treatment).

EXPERIMENT 3—RESPONSE OF EDAs AND EDVs TO EXOGENOUS CAPSAICIN

Concentration response curves to capsaicin^e (10⁻¹⁰ to 10⁻⁴M) were obtained for paired endothelium-intact and -denuded EDA and EDV from 9 horses. Responses of paired EDA(-e)s and EDV(-e)s (n = 6; 1 pair/horse) that had been precontracted with U44069 (3 \times 10⁻⁸M) were then obtained to a single concentration of h α CGRP (3 \times 10⁻⁷M), a single concentration of capsaicin (10⁻⁵M), and a vehicle control treatment (DMSO). The single concentrations of h α CGRP and capsaicin were chosen on the basis of the previous CRCs (experiment 1 and the first portion of experiment 3) in which a \leq 80% relaxant response in U44069-precontracted EDAs and EDVs had been elicited. The same volume of the vehicle control treatment (DMSO) was also added to additional vessels at basal resting tension for each horse.

EXPERIMENT 4—RELAXANT RESPONSES OF EDAs AND EDVs TO h α CGRP IN THE PRESENCE OF A SUBTHRESHOLD CONCENTRATION OF ET-1

Concentration response curves to h α CGRP (10⁻¹⁰ to 10⁻⁶M) were obtained for paired endothelium-intact and -denuded EDA and EDV from 6 horses in vessels precontracted with U44069 (3 \times 10⁻⁸M) following a 30-minute incubation with ET-1^e (10⁻¹²M) or the same volume of distilled water (vehicle control).

Statistical analysis—Data were expressed as decreases in tension (calculated as a percentage of the U44069-induced tension) and plotted against the log₁₀ CGRP concentration. Cumulative CRCs were fitted to a 1-site model by use of a computerized nonlinear curve-fitting program.^{f,g} The logistic equation used to fit the curves was as follows:

$$\text{Decrease in tension} = E_{\max} \cdot D^n / (D^n + EC_{50}^n)$$

where E_{max} represents the maximum response, D is the concentration of the drug used, n is the Hill slope (a measure of the gradient of the relationship between concentration and response) of the response curve, and EC₅₀ is the concentration of the drug at which the response was half of the maximum response. The EC₅₀ values were expressed as geometric means with 95% confidence intervals, whereas the E_{max} and Hill slope values were expressed as arithmetic means \pm SEM. The EC₅₀ and E_{max} values were compared by use of a paired Student *t* test^f or an ANOVA and Bonferroni *t* test,^h as appropriate. Significance was set at values of *P* \leq 0.05.

Results

Experiment 1—The CRC responses to h α CGRP were summarized (Table 1). The addition of h α CGRP induced a nonendothelium, concentration-dependent vasodilation in both EDAs and EDVs (Figure 1). Although there were no significant differences between the relaxant responses of the endothelial-intact and endothelial-denuded vessels to h α CGRP, the relaxant responses for both EDAs and EDVs to h α CGRP were consistently less in endothelial-denuded vessels. The relaxant response to h α CGRP was significantly greater in EDAs than in EDVs, both

Table 1—The CRC values for paired endothelium-intact and -denuded EDAs and EDVs (n = 6; 1 pair/horse) to hαCGRP.

Vessel type	EC ₅₀ (10 ⁻⁸ M)	E _{max} (%)	Hill slope
EDA(+e)	0.29 (-1.96 to 2.53)	159.3 ± 19.5*	2.48 ± 0.79
EDV(+e)	1.04 (-0.68 to 2.76)	70.0 ± 13.7	1.10 ± 0.16
EDA(-e)	2.30 (-4.81 to 9.42)	138.9 ± 32.6†	2.21 ± 0.93
EDV(-e)	2.29 (0.22 to 4.37)	37.1 ± 10.8	2.10 ± 0.56

Values for EC₅₀ are obtained at a hαCGRP concentration of 10⁻⁸M and are expressed as the geometric mean (95% confidence interval); values for E_{max} are expressed as the arithmetic mean ± SEM percentage of the U44609-induced tension; values for Hill slope are expressed as the arithmetic mean ± SEM.

*Value differs significantly (P ≤ 0.05; 2-way ANOVA and a Bonferroni t test) from the value for EDV(+e). †Value differs significantly (P ≤ 0.05; 2-way ANOVA and a Bonferroni t test) from the value for EDV(-e).

EC₅₀ = Concentration of the drug at which the response was half of the maximum response. E_{max} = Maximum response.




Figure 1—Concentration response curves of paired (n = 6; 1 pair/horse) EDA(+e)s and EDA(-e)s (A) and EDV(+e)s and EDV(-e)s (B) to hαCGRP. Results are expressed as the mean ± SEM percentage of the U44609-induced tension. In both panels, endothelial-intact and endothelial-denuded vessels are indicated by squares and triangles, respectively. Notice that the scales on the y-axis differ between panels.

with an intact endothelium (E_{max} value, 159.3 ± 19.5% of U44069-induced tension vs 70.0 ± 13.7% of U44069-induced tension, respectively; P = 0.02) and with a denuded endothelium (E_{max} value, 138.9 ± 32.6% of U44069-induced tension vs 37.1 ± 10.8% of U44069-induced tension, respectively; P = 0.05). There were no significant differences between the EC₅₀ and Hill slope values for any of the groups.




Figure 2—Responses of paired (n = 5; 1 pair/horse) EDA(-e)s (A) and EDV(-e)s (B) to a single concentration of hαCGRP (10⁻⁷M) before (black bar) and after (white bar) incubation with hαCGRP (10⁻⁷M) or the vehicle control treatment (1% acetic acid) for 30 minutes. Results are expressed as the mean ± SEM percentage of the U44609-induced tension. Notice that the scales on the y-axis differ between panels.

Experiment 2—The relaxation responses of EDA(-e) and EDV(-e) to hαCGRP before and after incubation with hαCGRP₈₋₃₇ or the vehicle control treatment did not differ significantly (Figure 2).

Experiment 3—In all vessels, capsaicin induced a concentration-dependent, biphasic relaxant CRC; there was a plateau in the response to capsaicin between concentrations of 3 × 10⁻⁸M and 3 × 10⁻⁶M (data not shown). The addition of a single concentration of hαCGRP or capsaicin to U44069-precontracted EDA(-e)s induced a strong relaxant response (E_{max} value, 103.5 ± 6.0% of U44069-induced tension vs 101.1 ± 1.7% of U44069-induced tension, respectively); both of these values differed significantly (P = 0.02) from the minimal relaxant response of U44069-precontracted EDA(-e)s to the vehicle control treatment (DMSO; Figure 3). The addition of a single concentration of hαCGRP or capsaicin to U44069-precontracted EDV(-e)s also induced a strong relaxant response (E_{max} value, 46.1 ± 10.2% of U44069-induced tension vs 47.3 ± 12.8% of U44069-induced tension, respectively); both of these values differed significantly (P = 0.02) from the relaxant response of U44069-precontracted EDV(-e)s to the vehicle control treatment (DMSO). There were minimal changes in basal resting tension in any of the vessel groups following the addition of DMSO.




Figure 3—Responses of paired ($n = 6$; 1 pair/horse) EDA(-e)s (A) and EDV(-e)s (B) to a single concentration of capsaicin (black bar), h α CGRP (gray bar), or the vehicle control treatment (DMSO [white bar]). Results are expressed as the mean \pm SEM percentage of the U44069-induced tension. Notice that the scales on the y-axis differ between panels. *Value for the maximum response differs significantly ($P < 0.001$) from the maximum response values for the other treatments.

Experiment 4—We did not detect significant differences between the relaxant responses to h α CGRP for EDA(+e) pretreated with ET-1 or the vehicle control treatment (E_{\max} value, $81.3 \pm 2.8\%$ of U44069-induced tension vs $74.3 \pm 7.4\%$ of U44069-induced tension, respectively) and for EDA(-e) pretreated with ET-1 or the vehicle control treatment (E_{\max} value, $78.7 \pm 2.1\%$ of U44069-induced tension vs $69.9 \pm 7.5\%$ of U44069-induced tension, respectively). There were no significant differences between the relaxant responses to h α CGRP for EDV(+e) incubated with or without ET-1 (E_{\max} value, $64.3 \pm 4.9\%$ of U44069-induced tension vs $61.2 \pm 3.5\%$ of U44069-induced tension, respectively) and for EDV(-e) pretreated with ET-1 or the vehicle control treatment (E_{\max} value, $48.9 \pm 6.5\%$ of U44069-induced tension vs $58.8 \pm 8.9\%$ of U44069-induced tension, respectively). There were no significant differences between the EC_{50} and Hill slope values for any of the groups.

Discussion

To our knowledge, in vitro responses of healthy equine digital vessels to CGRP have not been evaluated

previously. The relatively high potency and efficacy of h α CGRP as a vasodilator when exogenously applied to the digital vessels suggested that these responses are receptor-mediated events. Furthermore, the relaxant effects of h α CGRP on EDAs and EDVs were evident in the absence of the endothelium, which suggested that most of the receptors for CGRP are located on the vascular smooth muscle cells and that their activation leads to relaxation of the smooth muscle. However, the CRCs for EDA(+e) and EDV(+e) were to the left of the corresponding curves obtained for the EDA(-e) and EDV(-e) segments. This observation indicated that although the response to CGRP was not an endothelium-dependent event, mediators released from the endothelium may act synergistically with CGRP. It has been suggested that NO may mediate a portion of the vasorelaxant response of some vascular beds to CGRP.²⁴ However, the lack of synergistic effect of endothelial-derived relaxing factors and CGRP in equine digital vessels was supported by the fact that no significant differences were found between the EC_{50} and E_{\max} values for EDAs and EDVs with and without a functional endothelium in the present study. This lack of a significant difference indicated that the influence of the endothelium was variable, both across the range of concentrations tested and among horses, such that significant differences could not be detected with the number of horses included in the present study. Further investigations are necessary to determine the importance of tonic release of mediators (eg, NO, prostaglandin I₂, and endothelial-derived hyperpolarizing factor) from the endothelium on CGRP responses, perhaps by use of inhibitors of these mediators rather than physical removal of the endothelium that may damage the smooth muscle layer and concomitantly affect the direct relaxant effects of CGRP. However, in a preliminary study conducted by our research group on 3 horses (data not shown) in which the responses of paired endothelial-intact and -denuded arteries and veins were compared with responses for h α CGRP after incubation with a NO-synthase inhibitor (L-N^G-nitro-L-arginine methyl ester [300 μ M]) or a distilled water vehicle, no significant differences were revealed in the responses of the vessels to h α CGRP.

The in vitro responses of digital vessels to CGRP need to be characterized more fully before any firm conclusions can be drawn about the effects of this neuropeptide on equine digital vasculature. The high potency of CGRP suggests its effects are receptor-mediated events; however, the type of receptors involved and where these receptors are located have not been determined. The use of the selective peptide CGRP-receptor antagonist, h α CGRP₈₋₃₇, in the present study did not inhibit the relaxant responses of EDA(-e) or EDV(-e) to h α CGRP. The concentration of h α CGRP₈₋₃₇ used in the study reported here was selected on the basis of the pA₂ values reported for other species⁴⁵⁻⁵⁰ and on the basis of CRCs obtained for equine digital vessels in response to h α CGRP obtained in the present study. The single concentration of h α CGRP₈₋₃₇ (1 μ M) used in the present study was 10 times the reported affinity of this antagonist for CGRP-1 receptors in other species.⁴⁵⁻⁵⁰ It is possible that because α CGRP₈₋₃₇ has a higher affinity for the

CGRP-I receptor subtype ($pA_2 = 7$ to 8)⁴⁵⁻⁴⁸ than for the CGRP-II receptor subtype ($pA_2 = 5.5$ to 6.5),⁴⁸⁻⁵⁰ equine digital vasculature may contain primarily CGRP-II receptor subtypes. This is not considered likely because the responses of the digital vessels to low concentrations of $h\alpha$ CGRP were pronounced, which would make it more likely that the primary receptor in equine digital vasculature is the CGRP-I receptor subtype or an atypical CGRP-I receptor with an extremely low affinity for $h\alpha$ CGRP.⁸⁻³⁷ Interestingly, equine CGRP-I has low homology (51% to 59%)¹⁶ with CGRP-I from other species, whereas equine CGRP-II has greater homology (> 80%) with CGRPs from other species. Although the homology between human CGRP-I and human CGRP-II is high (> 90%), the homology between equine CGRP-I and equine CGRP-II is low (< 56%)¹⁶; thus, it has been speculated that CGRP-II in horses may have functions that are typically attributable to CGRP-I in other species.¹⁶

Although researchers have identified CGRP within the perivascular sensory-motor nerves in equine digits,³² its origin within the equine digital circulation and hence its function as an important endogenous vasodilator are unknown. Sensory-motor nerves can be classified as capsaicin-sensitive or capsaicin-insensitive, with capsaicin-sensitive nerves made up of primarily unmyelinated (C-fibers) and myelinated ($A\delta$ -fibers) nerve fibers that respond to chemical, thermal, and mechanical stimuli by releasing neurotransmitters such as CGRP.^{37,51,52} It is known that CGRP is primarily released from perivascular capsaicin-sensitive nerve endings,⁵³ although in rats, CGRP is also synthesized and stored within arterial endothelial cells.⁵⁴ Capsaicin, which acts on the vanilloid VR-1 receptor, can stimulate the release of CGRP (and SP) from capsaicin-sensitive nerves in humans,⁵⁵ guinea pigs,⁵⁶ and rats.⁵⁵ Thus, the in vitro vascular relaxant responses of endothelial-denuded vessels to exogenous $h\alpha$ CGRP and capsaicin observed in the present study supported the hypothesis that endogenous CGRP is released from perivascular sensory-motor nerves. However, the conclusion that capsaicin causes vasodilation through neuronal release of CGRP requires additional evidence for selective antagonists before it can be stated with certainty. In addition, electrical field stimulation in the presence of appropriate antagonists could provide further evidence that may support the hypothesis that endogenous CGRP is released from perivascular sensory-motor nerves.

Analysis of results of the experiments reported here indicated that $h\alpha$ CGRP is a potent and highly effective endothelium-independent vasodilator of both EDAs and EDVs, with significantly higher potency for EDAs. The possibility that relaxant factors released from the endothelium enhance the responses to CGRP warrants further investigation. The antagonist data were inconclusive, and additional functional and binding studies are required before the type and location of the receptors can be elucidated. Similar to the situation for SP, further studies are warranted to investigate the importance of CGRP-mediated vasodilation in protection against digital ischemia during acute laminitis in horses. On the basis of the vasodilatory responses of healthy equine digital vessels to the neuropeptides SP³⁹ and CGRP (observed in the study reported here), it is

likely that the peripheral nervous system is involved in the regulation of digital blood flow, with profound consequences when such regulation does not function properly.

- a. Vycon, Ecouen, France.
- b. Model HSE 30, isometric force transducer, Hugo Sachs Electronics, March-Hugstetten, Germany.
- c. Linseis 650 dual-channel pen recorder, Linton Instruments, Diss, Norfolk, England.
- d. Model HSE 301, bridge amplifier, Hugo Sachs Electronics, March-Hugstetten, Germany.
- e. Sigma Chemical Co, Poole, Dorset, England.
- f. GraphPad Prism, version 3.0 for Windows, GraphPad Software, San Diego, Calif.
- g. Bailey SR, Barlow RB, Elliott J, et al. SPLICE—a simple programme for the identification of curvaceous elements (abstr). *Br J Pharmacol* 1998;124:142P.
- h. SPSS, version 9.0 for Windows, SPSS Inc, Chicago, Ill.

References

1. Hunt RJ. The pathophysiology of acute laminitis. *Compend Contin Educ Pract Vet* 1991;13:401-407.
2. Moore JN, Allen D. The pathophysiology of acute laminitis. *Vet Med* 1996;91:936-939.
3. Burnstock G, Ralevic V. New insights into the local regulation of blood flow by perivascular nerves and endothelium. *Br J Plast Surg* 1994;47:527-543.
4. Coffman JR, Colles CM. Insulin tolerance in laminitic ponies. *Can J Comp Med* 1983;47:347-351.
5. Hood DM, Amoss MS, Grosenbaugh DA. Equine laminitis: a potential model of Raynaud's phenomenon. *Angiology* 1990;41:270-277.
6. Cerinic MM, Generini S, Pignone A. New approaches to the treatment of Raynaud's phenomenon. *Curr Opin Rheumatol* 1997;9:544-556.
7. Bunker CB, Terenghi G, Sprinall DR, et al. Deficiency of calcitonin gene-related peptide in Raynaud's phenomenon. *Lancet* 1990;336:1530-1533.
8. Jones E, Vinuela-Fernandez I, Eager RA, et al. Neuropathic changes in equine laminitis pain. *Pain* 2007;132:321-331.
9. Goltzman D, Mitchell J. Interaction of calcitonin and calcitonin gene-related peptide at receptor sites in target tissues. *Science* 1985;227:1343-1345.
10. Brain SD, Cambridge H. Calcitonin gene-related peptide: vasoactive effects and potential therapeutic role. *Gen Pharmacol* 1996;27:607-611.
11. Valentijn K, Gutow AP, Troiano N, et al. Effects of calcitonin gene-related peptide on bone turnover in ovariectomized rats. *Bone* 1997;21:269-274.
12. Feng Y, Tang Y, Guo J, et al. Inhibition of LPS-induced TNF- α production by calcitonin gene-related peptide (CGRP) in cultured mouse peritoneal macrophages. *Life Sci* 1997;61:PL281-PL287.
13. Gangula PR, Zhao H, Supowit SC, et al. Increased blood pressure in alpha-calcitonin gene-related peptide/calcitonin gene knockout mice. *Hypertension* 2000;35:470-475.
14. Monneret G, Pachot A, Laroche B, et al. Procalcitonin and calcitonin gene-related peptide decrease LPS-induced TNF production by human circulating blood cells. *Cytokine* 2000;12:762-764.
15. Amara SG, Arriza JL, Leff SE, et al. Expression in brain of a messenger RNA encoding a novel neuropeptide homologous to calcitonin gene-related peptide. *Science* 1985;229:1094-1097.
16. Toribio RE, Kohn CW, Leone GW, et al. Molecular cloning and expression of equine calcitonin, calcitonin gene-related peptide-I, and calcitonin gene-related peptide-II. *Mol Cell Endocrinol* 2003;199:119-128.
17. Poyner DR, Sexton PM, Marshall I, et al. International union of pharmacology. XXXII. The mammalian calcitonin gene-related peptides, adrenomedullin, amylin, and calcitonin receptors. *Pharmacol Rev* 2002;54:233-246.

18. Brain SD, Williams TJ, Tippins JR, et al. Calcitonin gene-related peptide is a potent vasodilator. *Nature* 1985;313:54–56.
19. DiPette DJ, Schwarzenberger K, Kerr N, et al. Dose dependent systemic and regional hemodynamic effects of calcitonin gene-related peptide. *Am J Med Sci* 1989;297:65–70.
20. Gibbins IL, Furness JB, Costa M. Pathway-specific patterns of the co-existence of substance P, calcitonin gene-related peptide, cholecystokinin and dynorphin in neurons of the dorsal root ganglia of the guinea-pig. *Cell Tissue Res* 1987;248:417–437.
21. Hanesch U, Heppelmann B, Schmidt RF. Substance P- and calcitonin gene-related peptide immunoreactivity in primary afferent neurons of the cat's knee joint. *Neuroscience* 1991;45:185–193.
22. Merighi A. Costorage and coexistence of neuropeptides in the mammalian CNS. *Prog Neurobiol* 2002;66:161–190.
23. Van Rossum D, Hanisch UK, Quirion R. Neuroanatomical localization, pharmacological characterization and functions of CGRP, related peptides and their receptors. *Neurosci Biobehav Rev* 1997;21:649–678.
24. Wimalawansa SJ. Calcitonin gene-related peptide and its receptors: molecular genetics, physiology, pathophysiology, and therapeutic potentials. *Endocr Rev* 1996;17:533–585.
25. Lundberg JM, Franco-Cereceda A, Hua X, et al. Co-existence of substance P and calcitonin gene-related peptide-like immunoreactivities in sensory nerves in relation to cardiovascular and bronchoconstrictor effects of capsaicin. *Eur J Pharmacol* 1985;108:315–319.
26. Sonea IM. Adrenergic and peptidergic innervation of the trachealis muscle in the normal horse: a preliminary report. *Res Vet Sci* 1993;54:335–339.
27. Sonea IM, Bowker RM, Robinson NE, et al. Substance P and calcitonin gene-related peptide-like immunoreactive nerve fibers in lungs from adult equids. *Am J Vet Res* 1994;55:1066–1074.
28. Sonea IM, Bowker RM, Robinson NE, et al. Distribution of SP- and CGRP-like immunoreactive nerve fibers in the lower respiratory tract of neonatal foals: evidence for loss during development. *Anat Embryol* 1994;190:469–477.
29. Corcoran BM, Mayhew IG, Hahn CN, et al. Expression of vasoactive intestinal peptide, calcitonin gene-related peptide, substance P, and intermediate neurofilaments in nasal mucosal nerve fibers of horses without nasal disease. *Am J Vet Res* 2000;61:1619–1624.
30. Yamamoto Y, Atoji Y, Hobo S, et al. Morphology of the nerve endings in laryngeal mucosa of the horse. *Equine Vet J* 2001;33:150–158.
31. Bae JE, Corcoran BM. Immunohistochemical study of the distribution of adrenergic and peptidergic innervation in the equine uterus and the cervix. *Reproduction* 2001;122:275–282.
32. Bowker RM, Brewer AM, Vex KB, et al. Sensory receptors in the equine foot. *Am J Vet Res* 1993;54:1840–1844.
33. Cornelissen BPM, Buma P, Rijkhuizen ABM, et al. Innervation of the equine mature and immature proximal sesamoid bone by calcitonin gene-related peptide and substance P-containing nerves. *Am J Vet Res* 1998;59:1378–1385.
34. Bowker RM, Rockershouser SJ, Vex KB, et al. Immunocytochemical and dye distribution studies of nerves potentially desensitized by injections into the distal interphalangeal joint or the navicular bursa of horses. *J Am Vet Med Assoc* 1993;203:1708–1714.
35. Bowker RM, Linder KE, Van Wulven KK, et al. Anatomy of the distal interphalangeal joint of the mature horse: relationships with navicular suspensory ligaments, sensory nerves and neurovascular bundle. *Equine Vet J* 1997;29:126–135.
36. Molyneux GS, Haller CJ, Mogg K, et al. The structure, innervation and location of arteriovenous anastomoses in the equine foot. *Equine Vet J* 1994;26:305–312.
37. Watson RE, Supowit SC, Zhao H, et al. Role of sensory nervous system vasoactive peptides in hypertension. *Braz J Med Biol Res* 2002;35:1033–1045.
38. Zhu F-X, Zhang X-Y, Olszewski MA, et al. Mechanism of capsaicin-induced relaxation in equine tracheal smooth muscle. *Am J Physiol* 1997;273:L997–L1001.
39. Katz LM, Marr CM, Elliott J. Characterisation of the response of equine digital arteries and veins to substance P. *J Vet Pharmacol Ther* 2003;26:361–368.
40. Miller RC, Mony M, Schini V, et al. Endothelial mediated inhibition of contraction and increase in GMP levels evoked by the alpha-adrenoreceptor agonist B-HT920 in rat isolated aorta. *Br J Pharmacol* 1984;83:903–908.
41. Elliott J, Bryant CE, Soydan J. The role of nitric oxide in the responses of equine digital veins to vasodilator and vasoconstrictor agents. *Equine Vet J* 1994;26:378–384.
42. Zerpa H, Bailey SR, Berhane Y, et al. Pharmacological characterization of alpha1-adrenoceptors in equine digital veins. *J Vet Pharmacol Ther* 2006;29:55–61.
43. Baxter GM, Laskey RE, Tackett RL, et al. In vitro reactivity of digital arteries and veins to vasoconstrictive mediators in healthy horses and in horses with early laminitis. *Am J Vet Res* 1989;50:508–517.
44. Bailey SR, Elliott J. Plasma 5-hydroxytryptamine constricts equine digital blood vessels in vitro: implications for pathogenesis of acute laminitis. *Equine Vet J* 1998;30:124–130.
45. Dennis TB, Fournier A, St Pierre S, et al. Structure-activity profile of calcitonin gene-related peptide in peripheral and brain tissues. Evidence for receptor multiplicity. *J Pharmacol Exp* 1989;251:718–725.
46. Dennis T, Fournier A, Cadieux A, et al. hCGRP₈₋₃₇, a calcitonin gene-related peptide antagonist revealing calcitonin gene-related peptide receptor heterogeneity in brain and periphery. *J Pharmacol Exp* 1990;254:123–128.
47. Quirion R, Van Rossum E, Dumont Y, et al. Characterization of CGRP1 and CGRP2 receptors. *Ann N Y Acad Sci* 1992;657:88–105.
48. Juaneda C, Dumont Y, Quirion R. The molecular pharmacology of CGRP and related peptide receptor subtypes. *Trends Pharmacol Sci* 2000;21:432–438.
49. Guiliani S, Wimalawansa SJ, Maggi CA. Involvement of multiple receptors in the biological effects of calcitonin gene-related peptide and amylin in rat and guinea-pig preparations. *Br J Pharmacol* 1992;107:510–514.
50. Longmore J, Hogg JE, Hutson PH, et al. Effects of two truncated forms of human calcitonin gene-related peptide: implications for receptor classification. *Eur J Pharmacol* 1994;265:53–59.
51. Holzer P. Local effector functions of capsaicin-sensitive sensory nerve endings: involvement of tachykinins, calcitonin gene-related peptide, and other neuropeptides. *Neurosci* 1988;24:739–768.
52. Rubino A, Burnstock G. Capsaicin-sensitive sensory-motor neurotransmission in the peripheral control of cardiovascular function. *Cardiovasc Res* 1996;31:467–479.
53. Zaidi M, Bevis PJ, Girgis SI, et al. Circulating CGRP comes from the perivascular nerves. *Eur J Pharmacol* 1985;117:283–284.
54. Doi Y, Kudo H, Nishino T, et al. Synthesis of calcitonin gene-related peptide (CGRP) by rat arterial endothelial cells. *Histol Histopathol* 2001;16:1073–1079.
55. Vaishnava P, Wang DH. Capsaicin sensitive-sensory nerves and blood pressure regulation. *Curr Med Chem Cardiovasc Hematol Agents* 2003;1:177–188.
56. Jansen I, Alafaci C, Uddman R, et al. Evidence that calcitonin gene-related peptide contributes to the capsaicin-induced relaxation of guinea pig cerebral arteries. *Regul Pept* 1993;31:167–178.