

- DB, ed. *Remington. The science and practice of pharmacy*. 21st ed. Baltimore: Lippincott Williams & Wilkins, 2006;861–862.
23. Featherstone HJ, Heinrich CL. Ophthalmic examination and diagnostics. In: Gelatt KN, Gilger BC, Kern TJ, eds. *Veterinary ophthalmology*. 5th ed. Ames, Iowa: Wiley-Blackwell, 2013;568–572.
 24. Haskins S, Pascoe PJ, Ilkiw JE, et al. Reference cardiopulmonary values in normal dogs. *Comp Med* 2005;55:156–161.
 25. Comline RS, Silver M. A comparative study of blood gas tensions, oxygen affinity and red cell 2,3 DPG concentrations in foetal and maternal blood in the mare, cow and sow. *J Physiol* 1974;242:805–826.
 26. Rhodes M, Heinrich C, Featherstone H, et al. Parotid duct transposition in dogs: a retrospective review of 92 eyes from 1999 to 2009. *Vet Ophthalmol* 2012;15:213–222.
 27. Lavy E, Goldberger D, Friedman M, et al. pH values and mineral content of saliva in different breeds of dogs. *Isr J Vet Med* 2012;67:244–248.
 28. Barrera R, Jimenez A, Lopez R, et al. Evaluation of total protein content in tears of dogs by polyacrylamide gel disk electrophoresis. *Am J Vet Res* 1992;53:454–456.
 29. Davidson HJ, Gerlach JA, Bull RW. Determination of protein concentrations and their molecular weight in tears from cats with normal corneas and cats with corneal sequestrum. *Am J Vet Res* 1992;53:1756–1759.
 30. Ollivier FJ. The precorneal tear film in horses: its importance and disorders. *Vet Clin North Am Equine Pract* 2004;20:301–318.
 31. Chen Z, Shamsi FA, Li K, et al. Comparison of camel tear proteins between summer and winter. *Mol Vis* 2011;17:323–331.



Correction: Ultrastructure and growth factor content of equine platelet-rich fibrin gels

In the report “Ultrastructure and growth factor content of equine platelet-rich fibrin gels” (*Am J Vet Res* 2014;75:392–401), information in the Materials and Methods is incorrect. On page 393, in the fourth sentence of the second paragraph under the heading Gel preparation, the amount of aprotinin added to each well to reduce proteolytic degradation is incorrect. The amount of aprotinin added to each well was 500 µg.