

- key eyes. I. Aqueous humor dynamics. *Invest Ophthalmol Vis Sci* 1987;28:463-469.
40. Tsai S, Miller PE, Struble C, et al. Topical application of 0.005% latanoprost increases episcleral venous pressure in normal dogs. *Vet Ophthalmol* 2012;15(suppl 1):71-78.
 41. Lindsey JD, Kashiwagi K, Boyle D, et al. Prostaglandins increase proMMP-1 and proMMP-3 secretion by human ciliary smooth muscle cells. *Curr Eye Res* 1996;15:869-875.
 42. Sagara T, Gatton DD, Lindsey JD, et al. Topical prostaglandin F2-alpha treatment reduces collagen types I, III, and IV in the monkey uveoscleral outflow pathway. *Arch Ophthalmol* 1999;117:794-801.
 43. Richter M, Krauss AHP, Woodward DF, et al. Morphological changes in the anterior eye segment after long-term treatment with different receptor selective prostaglandin agonists and a prostamide. *Invest Ophthalmol Vis Sci* 2003;44:4419-4426.
 44. Poyer JF, Millar C, Kaufman PL. Prostaglandin F2 alpha effects on isolated rhesus monkey ciliary muscle. *Invest Ophthalmol Vis Sci* 1995;36:2461-2465.
 45. Tofflemire KL, Whitley EM, Allbaugh RA, et al. Comparison of two- and three-times-daily topical ophthalmic application of 0.005% latanoprost solution in clinically normal dogs. *Am J Vet Res* 2015;76:625-631.
 46. Tsai S, Almazan A, Lee SS, et al. The effect of topical latanoprost on anterior segment anatomic relationships in normal dogs. *Vet Ophthalmol* 2013;16:370-376.
 47. Gelatt KN, MacKay EO. Effect of different dose schedules of latanoprost on intraocular pressure and pupil size in the glaucomatous Beagle. *Vet Ophthalmol* 2001;4:283-288.
 48. Park S, Kang S, Lee E, et al. Ultrasound biomicroscopic study of the effects of topical latanoprost on the anterior segment and ciliary body thickness in dogs. *Vet Ophthalmol* 2016;19:498-503.
-

Correction: Pharmacokinetics and pharmacodynamics of oral pradofloxacin administration in dogs

In the report "Pharmacokinetics and pharmacodynamics of oral pradofloxacin administration in dogs" (*Am J Vet Res* 2018;79:1268-1276), the recommended dose of pradofloxacin is incorrectly given in the final sentence of the structured abstract and final paragraph of the Discussion section as 7.5 mg/kg. The correct dose is 3 mg/kg. Under the Drugs heading in the Materials and Methods section, the approved dosage of pradofloxacin for dogs and cats in Europe is incorrectly given as 7.5 mg/kg, PO, every 24 hours. The correct dosage is 3 mg/kg, PO, every 24 hours.