

7. Soni A, Pepper GM, Wyrwinski PM, et al. Adrenal insufficiency occurring during septic shock: incidence, outcome, and relationship to peripheral cytokine levels. *Am J Med* 1995;98:266–271.
8. Moran JL, Chapman MJ, O'Fathartaigh MS, et al. Hypocortisolemia and adrenocortical responsiveness at onset of septic shock. *Intensive Care Med* 1994;20:489–495.
9. Feldman E, Nelson R. *Canine and feline endocrinology and reproduction*. 3rd ed. St Louis: Elsevier, 2004.
10. Cohn LA, Declue AE, Cohen RL, et al. Effects of fluticasone propionate dosage in an experimental model of feline asthma. *J Feline Med Surg* 2010;12:91–96.
11. Peterson ME, Greco DS, Orth DN. Primary hypoadrenocorticism in ten cats. *J Vet Intern Med* 1989;3:55–58.
12. Reinero CR, Brownlee L, Decile KC, et al. Inhaled flunisolide suppresses the hypothalamic-pituitary-adrenocortical axis, but has minimal systemic immune effects in healthy cats. *J Vet Intern Med* 2006;20:57–64.
13. Millard RP, Pickens EH, Wells KL. Excessive production of sex hormones in a cat with an adrenocortical tumor. *J Am Vet Med Assoc* 2009;234:505–508.
14. DeClue AE, Breshears LA, Pardo ID, et al. Hyperaldosteronism and hyperprogesteronism in a cat with an adrenal cortical carcinoma. *J Vet Intern Med* 2005;19:355–358.
15. Boag AK, Neiger R, Church DB. Trilostane treatment of bilateral adrenal enlargement and excessive sex steroid hormone production in a cat. *J Small Anim Pract* 2004;45:263–266.
16. Rossmeisl JH Jr, Scott-Moncrieff JC, Siems J, et al. Hyperadrenocorticism and hyperprogesteronemia in a cat with an adrenocortical adenocarcinoma. *J Am Anim Hosp Assoc* 2000;36:512–517.
17. Boord M, Griffin C. Progesterone secreting adrenal mass in a cat with clinical signs of hyperadrenocorticism. *J Am Vet Med Assoc* 1999;214:666–669.
18. Zimmer C, Horauf A, Reusch C. Ultrasonographic examination of the adrenal gland and evaluation of the hypophyseal-adrenal axis in 20 cats. *J Small Anim Pract* 2000;41:156–160.
19. Graves TK, Schall WD, Refsal K, et al. Basal and ACTH-stimulated plasma aldosterone concentrations are normal or increased in dogs with trichuriasis-associated pseudohypoadrenocorticism. *J Vet Intern Med* 1994;8:287–289.
20. Golden DL, Lothrop CD Jr. A retrospective study of aldosterone secretion in normal and adrenergic dogs. *J Vet Intern Med* 1988;2:121–125.
21. Willard MD, Refsal K, Thacker E. Evaluation of plasma aldosterone concentrations before and after ACTH administration in clinically normal dogs and in dogs with various diseases (Erratum published in *Am J Vet Res* 1988;49:283). *Am J Vet Res* 1987;48:1713–1718.
22. Goy-Thollot I, Decosne-Junot C, Bonnet JM. Influence of aging on adrenal responsiveness in a population of eleven healthy beagles. *Res Vet Sci* 2007;82:195–201.
23. Peterson ME, Kemppainen RJ. Dose-response relation between plasma concentrations of corticotropin and cortisol after administration of incremental doses of cosyntropin for corticotropin stimulation testing in cats. *Am J Vet Res* 1993;54:300–304.
24. Frank LA, Oliver JW. Comparison of serum cortisol concentrations in clinically normal dogs after administration of freshly reconstituted versus reconstituted and stored frozen cosyntropin. *J Am Vet Med Assoc* 1998;212:1569–1571.
25. Kemppainen R, Mansfield P, Sartin J. Endocrine responses of normal cats to TSH and synthetic ACTH administration. *J Am Anim Hosp Assoc* 1984;20:737–740.
26. Yu S, Morris JG. Plasma aldosterone concentration of cats. *Vet J* 1998;155:63–68.
27. Schoeman JP, Evans HJ, Childs D, et al. Cortisol response to two different doses of intravenous synthetic ACTH (tetracosactrin) in overweight cats. *J Small Anim Pract* 2000;41:552–557.
28. Tordjman K, Jaffe A, Trostanetsky Y, et al. Low-dose (1 microgram) adrenocorticotrophin (ACTH) stimulation as a screening test for impaired hypothalamo-pituitary-adrenal axis function: sensitivity, specificity and accuracy in comparison with the high-dose (250 microgram) test. *Clin Endocrinol (Oxf)* 2000;52:633–640.
29. Zaloga GP, Marik P. Hypothalamic-pituitary-adrenal insufficiency. *Crit Care Clin* 2001;17:25–41.
30. Peterson ME, Kemppainen RJ. Comparison of intravenous and intramuscular routes of administering cosyntropin for corticotropin stimulation testing in cats. *Am J Vet Res* 1992;53:1392–1395.
31. Peterson ME, Kemppainen RJ. Comparison of the immunoreactive plasma corticotropin and cortisol responses to two synthetic corticotropin preparations (tetracosactrin and cosyntropin) in healthy cats. *Am J Vet Res* 1992;53:1752–1755.
32. Pessina P, Fernández-Foren A, Cueto E, et al. Cortisol secretion after adrenocorticotrophin (ACTH) and dexamethasone tests in healthy female and male dogs. *Acta Vet Scand* 2009;51:33.
33. van Lier E, Perez-Clariget R, Forsberg M. Sex differences in cortisol secretion after administration of an ACTH analogue in sheep during the breeding and non-breeding season. *Anim Reprod Sci* 2003;79:81–92.
34. Javadi S, Slingerland LI, van de Beek MG, et al. Plasma renin activity and plasma concentrations of aldosterone, cortisol, adrenocorticotrophic hormone, and alpha-melanocyte-stimulating hormone in healthy cats. *J Vet Intern Med* 2004;18:625–631.
35. Willemsse T, Vroom MW, Mol JA, et al. Changes in plasma cortisol, corticotropin, and  $\alpha$ -melanocyte-stimulating hormone concentrations in cats before and after physical restraint and intradermal testing. *Am J Vet Res* 1993;54:69–72.



Correction: Effects of acepromazine, hydromorphone, or an acepromazine-hydromorphone combination on the degree of sedation in clinically normal dogs

In the article “Effects of acepromazine, hydromorphone, or an acepromazine-hydromorphone combination on the degree of sedation in clinically normal dogs” (*J Am Vet Med Assoc* 2010;237:1155–1159), the dosage of acepromazine was incorrect. The correct dosage of acepromazine is 0.05 mg/kg (0.023 mg/lb).