

4. Aikawa T, Fujita H, Kanazono S, et al. Long-term neurologic outcome of hemilaminectomy and disk fenestration for treatment of dogs with thoracolumbar intervertebral disk herniation: 831 cases (2000–2007). *J Am Vet Med Assoc* 2012;241:1617–1626.
5. Packer RMA, Hendricks A, Volk HA, et al. How long and low can you go? Effect of conformation on the risk of thoracolumbar intervertebral disc extrusion in domestic dogs. *PLoS ONE* 2013;8:e69650.
6. Goggin JE, Li AS, Franti CE. Canine intervertebral disk disease characterization by age, sex, breed, and anatomic site of involvement. *Am J Vet Res* 1970;31:1687–1692.
7. Priester WA. Canine intervertebral disc disease—occurrence by age, breed, and sex among 8,117 cases. *Theriogenology* 1976;6:293–303.
8. Brown NO, Helphrey ML, Prata RG. Thoracolumbar disk disease in dogs: a retrospective analysis of 187 cases. *J Am Anim Hosp Assoc* 1977;13:665–672.
9. Olby N, Levine J, Harris T, et al. Long-term functional outcome of dogs with severe injuries of the thoracolumbar spinal cord: 87 cases (1996–2001). *J Am Vet Med Assoc* 2003;222:762–769.
10. Ruddle TL, Allen DA, Schertel ER, et al. Outcome and prognostic factors in non-ambulatory Hansen type I intervertebral disc extrusions: 308 cases. *Vet Comp Orthop Traumatol* 2006;19:29–34.
11. Aikawa T, Fujita H, Shibata M, et al. Recurrent thoracolumbar intervertebral disc extrusion after hemilaminectomy and concomitant prophylactic fenestration in 662 chondrodystrophic dogs. *Vet Surg* 2012;41:381–390.
12. Roach WJ, Thomas M, Weh M, et al. Residual herniated disc material following hemilaminectomy in chondrodystrophic dogs with thoracolumbar intervertebral disc disease. *Vet Comp Orthop Traumatol* 2012;25:109–115.
13. Ferreira AJA, Correia JHD, Jaggy A. Thoracolumbar disc disease in 71 paraplegic dogs: influence of rate of onset and duration of clinical signs on treatment results. *J Small Anim Pract* 2002;43:158–163.
14. Ryan TM, Platt SR, Llabres-Diaz FJ, et al. Detection of spinal cord compression in dogs with cervical intervertebral disc disease by magnetic resonance imaging. *Vet Rec* 2008;163:11–15.
15. Dewey CW. Disorders of the cauda equina. In: *A practical guide to canine and feline neurology*. 2nd ed. Ames, Iowa: Wiley-Blackwell, 2008;389–404.
16. Levine GJ, Levine JM, Budke CM, et al. Description and repeatability of a newly developed spinal cord injury scale for dogs. *Prev Vet Med* 2009;89:121–127.
17. Van Wie EY, Fosgate GT, Mankin JM, et al. Prospectively recorded versus medical record-derived spinal cord injury scores in dogs with intervertebral disk herniation. *J Vet Intern Med* 2013;27:1273–1277.
18. Olby N, Harris T, Burr J, et al. Recovery of pelvic limb function in dogs following acute intervertebral disk herniations. *J Neurotrauma* 2004;21:49–59.
19. De Decker S, Wawrzynski LA, Volk HA. Clinical signs and outcome of dogs treated medically for degenerative lumbosacral stenosis: 98 cases (2004–2012). *J Am Vet Med Assoc* 2014;245:408–413.
20. Rohdin C, Jeserevic J, Viitmaa R, et al. Prevalence of radiographic detectable intervertebral disc calcifications in Dachshunds surgically treated for disc extrusion. *Acta Vet Scand* 2010;52:24.
21. Aikawa T, Shibata M, Asano M, et al. A comparison of thoracolumbar intervertebral disc extrusion in French Bulldogs and Dachshunds and association with congenital vertebral anomalies. *Vet Surg* 2014;43:301–307.
22. Smolders LA, Bergknut N, Grinwis GCM, et al. Intervertebral disc degeneration in the dog. Part 2: Chondrodystrophic and non-chondrodystrophic breeds. *Vet J* 2013;195:292–299.
23. Parker HG, Von Holdt BM, Quignon P, et al. An expressed *fgf4* retrogene is associated with breed-defining chondrodysplasia in domestic dogs. *Science* 2009;325:995–998.
24. Mogensen MS, Karlskov-Mortensen P, Proschowsky HF, et al. Genome-wide association study in dachshund: identification of a major locus affecting intervertebral disc calcification. *J Hered* 2011;102(suppl 1):S81–S86.
25. Mogensen MS, Scheibye-Alsing K, Karlskov-Mortensen P, et al. Validation of genome-wide intervertebral disc calcification associations in Sachshund and further investigation of the chromosome 12 susceptibility locus. *Front Genet* [serial online] 2012;3:225 Available at: journal.frontiersin.org/article/10.3389/fgene.2012.00225/full. Accessed Dec 17, 2014.
26. Braund KG, Ghosh P, Taylor TK, et al. Morphological studies of the canine intervertebral disc. The assignment of the Beagle to the achondroplastic classification. *Res Vet Sci* 1975;19:167–172.
27. Forterre F, Konar M, Spreng D, et al. Influence of intervertebral disc fenestration at the herniation site in association with hemilaminectomy on recurrence in chondrodystrophic dogs with thoracolumbar disc disease: a prospective MRI study. *Vet Surg* 2008;37:399–405.
28. Bergknut N, Auriemma E, Wijsman S, et al. Evaluation of intervertebral disc degeneration in chondrodystrophic and non-chondrodystrophic dogs by use of Pfirrmann grading of images obtained with low-field magnetic resonance imaging. *Am J Vet Res* 2011;72:893–898.
29. Dhupa S, Glickman NW, Waters DJ. Functional outcome in dogs after surgical treatment of caudal lumbar intervertebral disc herniation. *J Am Anim Hosp Assoc* 1999;35:323–331.
30. De Lahunta A, Glass E. Lower motor neuron: spinal nerve, general somatic efferent system. In: *Veterinary neuroanatomy and clinical neurology*. 3rd ed. St Louis: Elsevier Saunders, 2009;77–133.
31. Karli P, Martlé V, Bossens K, et al. Dominance of chemokine ligand 2 and matrix metalloproteinase-2 and -9 and suppression of pro-inflammatory cytokines in the epidural compartment after intervertebral disc extrusion in a canine model. *Spine J* 2014;14:2976–2984.
32. Kobayashi S, Takeno K, Miyazaki T, et al. Effects of arterial ischemia and venous congestion on the lumbar nerve root in dogs. *J Orthop Res* 2008;26:1533–1540.
33. Kobayashi S. Pathophysiology, diagnosis and treatment of intermittent claudication in patients with lumbar canal stenosis. *World J Orthop* 2014;5:134–145.
34. Gödde T, Steffen F. Surgical treatment of lumbosacral foraminal stenosis using a lateral approach in twenty dogs with degenerative lumbosacral stenosis. *Vet Surg* 2007;36:705–713.
35. Kubicek L, Kopáček R, Klusáková I, et al. Alterations in the vascular architecture of the dorsal root ganglia in a rat neuropathic pain model. *Ann Anat* 2010;192:101–106.



## Correction: Pathology in Practice: canine distemper virus infection in a puppy

In the Pathology in Practice report describing canine distemper virus (CDV) infection in a puppy (*J Am Vet Med Assoc* 2015;247:1375–1377), the sentence at the end of the first paragraph in the Comments section lists cheetahs as one of the many varieties of carnivores susceptible to CDV infection. However, it is not clear that cheetahs are in fact susceptible. For example, Appel et al<sup>1</sup> tested serum samples collected from 65 cheetahs between 1985 and 1993 at the National Zoo in Washington, DC, for antibodies against CDV, and results were negative for all 65. Munson et al<sup>2</sup> identified anti-CDV antibodies in serum samples from wild Namibian cheetahs; however, clinical disease was not documented, and the Association of Zoos and Aquariums Species Survival Plan does not currently recommend vaccinating cheetahs against CDV.

1. Appel MJ, Yates RA, Goley GL, et al. Canine distemper epizootic in lions, tigers, and leopards in North America. *J Vet Diagn Invest* 1994;6:277–288.
2. Munson L, Marker L, Dubovi E, et al. Serosurvey of viral infections in free-ranging Namibian cheetahs (*Acinonyx jubatus*). *J Wildl Dis* 2004;40:23–31.