

Supplementary Table S1—Minimal inhibitory concentration ($\mu\text{g/ml}$) profiles of the two *Staphylococcus aureus* strains (ATCC 29213; ATCC 43300) used in a study to assess the effects of autologous platelet-rich plasma on *Staphylococcus aureus*–induced chondrocyte death in an in vitro model of bovine septic arthritis.

Antimicrobial	<i>S aureus</i> ATCC Strain	
	29213	43300
Ampicillin	1 (R)*	1 (R)
Ceftriaxone	≤ 8 (S)	≤ 8 (R)
Ciprofloxacin	≤ 0.5 (S)	≤ 0.5 (S)
Clindamycin	≤ 0.12 (S)	≤ 0.12 (S)
Daptomycin	≤ 0.25 (S)	≤ 0.25 (S)
Erythromycin	≤ 0.25 (S)	≤ 0.25 (S)
Gatifloxacin	≤ 1 (NI)	≤ 1 (NI)
Gentamicin	≤ 2 (S)	≤ 2 (S)
Levofloxacin	≤ 0.25 (S)	≤ 0.25 (S)
Linezolid	2 (S)	2 (S)
Oxacillin + 2% NaCl	≤ 0.25 (S)	> 8 (R)
Penicillin	1 (R)	1 (R)
Quinupristin/dalfopristin	0.25 (S)	0.5 (NI)
Rifampin	≤ 0.5 (S)	≤ 0.5 (S)
Tetracycline	≤ 2 (S)	≤ 2 (S)
Trimethoprim/sulfamethoxazole	≤ 0.5 (S)	≤ 0.5 (S)
Vancomycin	≤ 1 (S)	≤ 1 (S)

(S) indicates susceptible, (R) indicates resistant, and (NI) indicates no interpretation based on CLSI breakpoint guidelines.