

Supplemental Table S1. Proportion of culture positive samples by affected anatomical structure

	% Culture Positive (n)	95% Confidence Interval
Overall	37.4 (951)	34.4, 40.6
<b>Affected Anatomical Structure</b>		
Bursa	52.1 (48)	38.3, 65.5
Bicipital	80 (5)	37.6, 96.4
Calcaneal	52.9 (17)	30.9, 73.8
Supracarpal	100 (1)	20.7, 100
Cubital	50 (2)	9.5, 90.5
Cunean	0 (1)	
Infraspinatous	100 (1)	20.7, 100
Navicular	60 (15)	35.7, 80.1
Nuchal	0 (5)	
Unknown Bursa	0 (1)	
Tendon Sheath	40.2 (72)	29.7, 51.8
Carpal Canal	33.3 (7)	8.2, 64.1
Common Digital Extensor	100 (1)	20.7, 100
Digital Flexor	37.3 (59)	26.1, 50
Extensor Carpi Radialis	100 (2)	34.2, 100
Long Digital Extensor	50 (2)	9.5, 90.5
Tarsal	100 (1)	20.7, 100
Joint	29.4 (482)	25.6, 33.7
Carpal	20.8 (72)	13.1, 31.6
Distal Interphalangeal	42.2 (45)	29.0, 56.7
Radiohumeral	46.7 (15)	24.8, 69.9
Metacarpo-/metatarsophalangeal	20.9 (91)	13.8, 30.3
Coxofemoral	83.3 (6)	43.6, 97
Tarsal	27.2 (184)	21.3, 34
Proximal Interphalangeal	33.3 (6)	9.7, 70
Glenohumeral	50 (6)	18.8, 81.2
Intersternal	0 (1)	
Stifle	39.2 (51)	27, 52.9
Temporomandibular	100 (1)	20.7, 100
Atlanto-occipital	100 (1)	20.7, 100
Intervertebral	0 (2)	
Unknown Joint	0 (1)	
Total	32.6 (598)	