

Supplementary Table S5. Tables illustrating comparisons of supernatant cytokine concentrations (pg/mL) in whole blood incubated for 12 h versus 24 h. Five individual tables based on rCTS1₁₀₅₋₃₁₀ antigen concentration (i.e., 10 µg/mL, 25 µg/mL, 50 µg/mL, 75 µg/mL) or vehicle control [phosphate-buffered saline (PBS)]. Data was analyzed using Wilcoxon signed-rank tests and are presented as mean and standard deviation

PBS Control

	12 h (n = 6)	24 h (n = 6)	p-value
IL-1β			
Mean (SD)	102.880 (0.000)	110.500 (18.665)	0.40
Range	102.880 - 102.880	102.880 - 148.600	
IL-2			
Mean (SD)	47.107 (14.591)	51.313 (24.895)	1.00
Range	41.150 - 76.890	41.150 - 102.130	
IL-12			
Mean (SD)	118.260 (57.096)	119.398 (44.697)	0.52
Range	41.150 - 216.540	41.150 - 165.810	
IL-4			
Mean (SD)	296.843 (213.658)	352.292 (299.073)	0.73
Range	205.760 - 732.860	205.760 - 959.420	
IL-6			
Mean (SD)	239.005 (81.433)	269.592 (135.541)	0.60
Range	205.760 - 405.230	205.760 - 543.820	
IL-8			
Mean (SD)	442.323 (637.907)	913.337 (1027.100)	0.23
Range	41.150 - 1647.000	90.600 - 2773.000	
MCP-1			
Mean (SD)	2277.333 (1053.958)	2438.833 (1237.086)	0.75
Range	1235.000 - 4237.000	1235.000 - 4760.000	
TNF-α			
Mean (SD)	107.218 (76.532)	112.988 (106.986)	0.75
Range	41.150 - 246.690	41.150 - 322.920	