

Supplemental Appendix S1—Anesthetic drugs included in the anesthesia care plan for a 6-month-old female French Bulldog undergoing balloon valvuloplasty for treatment of pulmonic stenosis. Anesthesia care occurred via real-time remote telecommunication provided by an experienced, board-certified anesthesiologist located in Liege, Belgium, communicating with the local anesthesiologist treating the patient throughout surgery taking place in Jelgava, Latvia. Communication (in English) occurred via an internet video communication application that conveyed real-time views of the patient, anesthesia machine, and monitors to the remote anesthesiologist. Names and doses of all drugs were verified and compared by the remote and local anesthesiologists during preanesthetic planning.

Drug	Belgium	Latvia
Fentanyl	Fentanyl (Janssen-Cilag) (0.5 mg/10 mL)	Fentanyl citrate-Kalceks (0.05 mg/mL)
Midazolam	Dormicum (15 mg/3 mL)	Fulsed (5 mg/mL)
Propofol	Diprivan (200 mg/20 mL)	Claris propofol (200 mg/20 ml)
Lidocaine	Linisol (200 mg/10 mL)	Lidocaine Claris (100 mg/5 mL)

Copyright 2017, American Veterinary Medical Association

Anesthesia Case of the Month. Tutunaru, AC; Chervensky, R; Ozolins, V; et al. *J Am Vet Med Assoc* 2017;250:984-987.

Available at: <http://avmajournals.avma.org/toc/javma/250/9>