

**Supplementary Table S2—Changes in clinical severity scores after endoluminal tracheal stent placement.**

**Percent of dogs change in clinical score\* compared to preoperative score**

Clinical sign	Short-term			Intermediate-term			Long-term			Final follow-up		
	Improved	Stable	Worsened	Improved	Stable	Worsened	Improved	Stable	Worsened	Improved	Stable	Worsened
Coughing	47 (18)	21 (8)	32 (12)	48 (13)	8 (2)	44 (12)	44 (15)	3 (1)	53 (18)	43 (22)	16 (8)	41 (21)
Goose-honking or raspy breathing	<b>94 (33)</b>	<b>3 (1)</b>	<b>3 (1)</b>	<b>92 (25)</b>	<b>4 (1)</b>	<b>4 (1)</b>	<b>88 (28)</b>	<b>3 (1)</b>	<b>9 (3)</b>	<b>89 (42)</b>	<b>2 (1)</b>	<b>9 (4)</b>
Dyspnea	<b>87 (32)</b>	<b>8 (3)</b>	<b>5 (2)</b>	<b>85 (24)</b>	<b>4 (1)</b>	<b>11 (3)</b>	<b>89 (31)</b>	<b>0 (0)</b>	<b>11 (4)</b>	<b>84 (43)</b>	<b>10 (5)</b>	<b>6 (3)</b>

Data represent percent of dogs (number of dogs reported). Increase in score indicated a worsening of clinical signs, decrease indicated an improvement, and stable indicated no change. Values in bold indicate the average clinical score for that period was statistically improved compared to the average preoperative score.

Refer to Table 2 for remainder of key.