

Supplementary Table S4—Comparison of long-term complications between standard incisional gastropexy (SIG) and modified incisional gastropexy (MIG) in each of six matched subgroups. Some cases are not included because they did not fit in any of the matched subgroups.

Subgroupings		No complication	Gastric dilatation	Trouble eating	Seroma	Unknown
1	SIG for GDV (n=91)	5	1	2	0	83
	MIG for GDV (n=16)	9	0	0	0	7
2	Prophylactic SIG (n=21)	2	0	0	0	19
	Prophylactic MIG (n=1)	1	0	0	0	0
3	OHE with SIG (n=71)	11	1	0	1	58
	OHE with MIG (n=12)	8	1	0	0	3
4	Castration with SIG (n=21)	1	0	0	0	20
	Castration with MIG (n=2)	1	0	0	0	1
5	Splenectomy with SIG (n=55)	1	0	0	0	54
	Splenectomy with MIG (n=6)	6	0	0	0	0
6	Celiotomy with SIG (n=12)	1	0	0	0	11
	Celiotomy with MIG (n=2)	1	0	0	0	1

Gastric dilatation = dilatation without volvulus

Unknown = no follow-up at this time; therefore, unable to determine if complications occurred

GDV = gastric dilatation-volvulus

OHE = ovariohysterectomy

Celiotomy = abdominal procedures other than GDV, OHE, or splenectomy